arXiv:1505.06622v2 [cs.LO] 16 Apr 2016

MONADIC SECOND ORDER FINITE SATISFIABILITY AND UNBOUNDED TREE-WIDTH

TOMER KOTEK, HELMUT VEITH, FLORIAN ZULEGER

Abstract. The finite satisfiability problem of monadic second order logic is decidable only on classes of structures of bounded tree-width by the classic result of Seese [24]. We prove that the following problem is decidable:

Input: (i) A monadic second order logic sentence α, and (ii) a sentence β in the two-variable fragment of first order logic extended with counting quantifiers. The vocabularies of α and β may intersect.

Output: Is there a finite structure which satisfies $\alpha \wedge \beta$ such that the restriction of the structure to the vocabulary of α has bounded tree-width? (The tree-width of the desired structure is not bounded.)

As a consequence, we prove the decidability of the satisfiability problem by a finite structure of bounded tree-width of a logic $\mathrm{MSO}^{\exists\mathrm{card}}$ extending monadic second order logic with linear cardinality constraints of the form $|X_1| + \cdots + |X_r| < |Y_1| + \cdots + |Y_s|$ on the variables X_i, Y_j of the outer-most quantifier block. We prove the decidability of a similar extension of WS1S.

1. Introduction

Monadic second order logic (MSO) is among the most expressive logics with good algorithmic properties. It has found countless applications in computer science in diverse areas ranging from verification and automata theory [13, 18, 25] to combinatorics [16, 17], and parameterized complexity theory [8, 6].

The power of MSO is most visible over graphs of bounded tree-width, and with second order quantifiers ranging over sets of edges[1]: (1) Courcelle's famous theorem shows that MSO model checking is decidable over graphs of bounded tree-width in linear time [5, 1]. (2) Finite satisfiability by graphs of bounded tree-width is decidable [5] (with non-elementary complexity) – thus contrasting Trakhtenbrot's undecidability result of first order logic. (3) Seese proved [24] that for each class $\mathscr{K}$ of graphs with *unbounded* tree-width, finite satisfiability of MSO by graphs in $\mathscr{K}$ is undecidable. Together, (2) and (3) give a fairly clear picture of the decidability of finite satisfiability of MSO. It appeared that (3) gives a natural limit for decidability of MSO on graph classes. For instance, finite satisfiability on planar graphs is undecidable because their tree-width is unbounded.

While Courcelle and Seese circumvent Trakhtenbrot's undecidability result by *restricting the classes of graphs (or relational structures)*, several other research communities have studied syntactic restrictions of first order logic. Modal logic [26], many temporal logics [21], [23, Chapter 24], the guarded fragment [9], many description logics [2], and the two-variable fragment [10] are restricted first order logics with decidable finite satisfiability, and hundreds of papers on these topics have explored the border between decidability and undecidability. While many of the earlier papers exploited variations of the tree model property to show decidability, recent research has also focused on logics such as the two-variable fragment with counting C^2 [11, 22], where finite satisfiability is decidable despite the absence of the tree model property. In a recent breakthrough result, Charatonik and Witkowski [4] have extended this result to structures with built-in binary trees. Note that this logic is not a fragment of first order logic, but more naturally understood as a very weak second order logic which can express one specific second order property – the property of being a tree.

---

[1] The logic we denote by MSO is denoted MS_2 by Courcelle and Engelfriet [6].

Our main result is a powerful generalization of the seminal result on decidability of the satisfiability problem of MSO over bounded tree-width and the recent theorem by [4]: We show decidability of finite satisfiability of conjunctions $\alpha \wedge \beta$ where α is in MSO and β is in C^2 by a finite structure $\mathfrak{M}$ whose restriction to the vocabulary of α has bounded tree-width. (Theorem 1 in Section 3)

Let us put this result into perspective:

- The MSO decidability problem is a trivial consequence by setting β to true; Charatonik and Witkowski's result follows by choosing α to be an MSO formula which axiomatizes a d-ary tree, which is a standard construction [6].
- The decidability of *model checking* $\alpha \wedge \beta$ over a finite structure is a much simpler problem than ours: We just have to model check α and β one after the other. In contrast, satisfiability is not obvious because α and β can share relational variables. running two finite satisfiability algorithms for the two formulas independently may tield two models which disagree on the shared vocabulary. Thus, the problem we consider is similar in spirit to (but technically very different from) Nelson-Oppen [20] combinations of theories.
- Our result trivially generalizes to Boolean combinations of sentences in the two logics.

Proof Technique. We show how to reduce our satisfiability problem for $\alpha \wedge \beta$ to the finite satisfiability of a C^2-sentence with a built-in tree, which is decidable by [4]. The most significant technical challenge is to eliminate shared binary relation symbols between α' and β. Our Separation Theorem overcomes this challenge by an elegant construction based on local types of universe elements and a coloring argument for directed graphs. The second technical challenge is to replace the MSO-sentence α with an equi-satisfiable C^2-sentence α'. To do so, we apply tools including the Feferman-Vaught theorem for MSO and translation schemes.

Monadic Second Order Logic with Cardinalities. Our main theorem imply new decidability results for monadic second order logic with cardinality constraints, i.e., expressions of the form $|X_1| + \ldots |X_r| < |Y_1| + \ldots |Y_t|$ where the X_i and Y_i are monadic second order variables. Klaedtke and Rueß [15] showed that the decision problem for the theory of weak monadic second order logic with cardinality constraints of one successor (WS1S$^{\mathrm{card}}$) is undecidable; they describe a decidable fragment where the second order quantifiers have no alternation and appear *after* the first order quantifiers in the prefix. Our main theorem implies decidability of a different fragment of WS1S with cardinalities: The fragment MSO$^{\exists\mathrm{card}}$ consists of formulas $\exists \bar{X} \psi$ where the cardinality constraints in ψ involve only the monadic



(a) (b)

Figure 1.

second order variables from $\bar{X}$, cf. Theorem 6 in Section 7. Note that in contrast to [15], our fragment is a strict superset of WS1S.

For WS2S, we are not aware of results about decidable fragments with cardinalities. We describe a strict superset of MSO whose satisfiability problem over finite graphs of bounded tree-width is decidable, and which is syntactically similar to the WS1S extension above.

Expressive Power over Structures. Our main result extends the existing body of results on finite satisfiability by structures of bounded tree-width to a significantly richer set of structures. The structures we consider are C^2-axiomatizable extensions of structures of bounded tree-width. For instance, we can have interconnected doubly-linked lists as in Fig. 1(a), or a tree whose leaves are connected in a chain and have edges pointing to any of the nodes of a cyclic list as in Fig. 1(b). Such structures occur very naturally as shapes of dynamic data structures in programming – where cycles and trees are containers for data, and additional edges express relational information between the data. The analysis of semantic relations between data structures served as a motivation for us to investigate the logics in the current paper [3].

Being a cyclic list or a tree whose leaves are chained can be expressed in MSO and both of these data structures have tree-width at most 3. We can compel the edges between the tree and the cyclic list to obey C^2-expressible constraints such as:

- every leaf of the tree has a single edge to the cyclic list;
- every node of the cyclic list has an incoming edge from at least one leaf of the tree; or
- any two leaves pointing to the same node of the cyclic list agree on membership in some unary relation.

Note that while the structures we consider may contain grids of unbounded sizes as subgraphs, the logic cannot axiomatize them.

2. Background

This section introduces basic definitions and results in model theory and graph theory. We follow [19] and [6].

The **two-variable fragment with counting** C^2 is the extension of the two-variable fragment of first order logic with first order counting quantifiers $\exists^{\leq n}$, $\exists^{\geq n}$, $\exists^{=n}$, for every $n \in \mathbb{N}$. Note that C^2 remains a fragment of first order logic.


(a) (b)

FIGURE 2.

Monadic Second order logic MSO is the extension of first order logic with set quantifiers which can range over elements of the universe or subsets of relations[2]. Throughout the paper all structures consist of unary and binary relations only. Structures are finite unless explicitly stated otherwise (in the discussion of WS1S). Let $\mathcal{C}$ be a vocabulary (signature). The **arity** of a relation symbol $C \in \mathcal{C}$ is denoted by $arity(C)$. The set of unary (binary) relation symbols in $\mathcal{C}$ are $\mathrm{un}(\mathcal{C})$ $(\mathrm{bin}(\mathcal{C}))$. We write $\mathrm{MSO}(\mathcal{C})$ for the set of MSO-formulas on the vocabulary $\mathcal{C}$. The quantifier rank of a formula $\varphi \in \mathrm{MSO}$, i.e. the maximal depth of nested quantifiers in φ is denoted $\mathrm{qr}(\varphi)$. We denote by $\mathfrak{A}_1 \sqcup \mathfrak{A}_2$ the **disjoint union** of two $\mathcal{C}$-structures $\mathfrak{A}_1$ and $\mathfrak{A}_2$. Given vocabularies $\mathcal{C}_1 \subseteq \mathcal{C}_2$, a $\mathcal{C}_2$-structure $\mathfrak{A}_2$ is an **expansion** of a $\mathcal{C}_1$-structure $\mathfrak{A}_1$ if $\mathfrak{A}_1$ and $\mathfrak{A}_2$ agree on the symbols in $\mathcal{C}_1$; in this case $\mathfrak{A}_1$ is the **reduct** of $\mathfrak{A}_2$ to $\mathcal{C}_1$, i.e. $\mathfrak{A}_1$ is the $\mathcal{C}_1$-reduct of $\mathfrak{A}_2$. We denote the reduct of $\mathfrak{A}_2$ to $\mathcal{C}_1$ by $\mathfrak{A}_2|_{\mathcal{C}_1}$. A $\mathcal{C}$-structure $\mathfrak{A}_0$ with universe A_0 is a **substructure** of a $\mathcal{C}$-structure $\mathfrak{A}_1$ with universe A_1 if $A_0 \subseteq A_1$ and for every $R \in \mathcal{C}$, $R^{\mathfrak{A}_0} = R^{\mathfrak{A}_1} \cap A_0^{arity(R)}$. We say that $\mathfrak{A}_0$ is the substructure of $\mathfrak{A}_1$ generated by A_0.

Graphs are structures of the vocabulary[3] $\mathcal{C}_G = \langle s \rangle$ consisting of a single binary relation symbol s. Graphs are simple and undirected unless explicitly stated otherwise. **Tree-width** $\mathrm{tw}(G)$ is a graph parameter indicating how close a simple undirected graph G is to being a tree, cf. [6]. It is well-known that a graph has tree-width at most k iff it is a partial k-tree. A **partial** k**-tree** is a subgraph of a k-tree. k**-trees** are built inductively from the $(k+1)$-clique by repeated addition of vertices, each of which is connected with k edges to a k-clique. The **Gaifman graph** $\mathrm{Gaif}(\mathfrak{A})$ of a $\mathcal{C}$-structure $\mathfrak{A}$ is the graph whose vertex set is the universe of $\mathfrak{A}$ and whose edge set is the union of the symmetric closures of $C^{\mathfrak{A}}$ for every $C \in \mathrm{bin}(\mathcal{C})$. Note the unary relations of $\mathfrak{A}$ play no role in $\mathrm{Gaif}(\mathfrak{A})$. The **tree-width** $\mathrm{tw}(\mathfrak{A})$ **of a** $\mathcal{C}$**-structure** $\mathfrak{A}$ is the tree-width of its Gaifman graph. In this paper, tree-width is a parameter of *finite* structures only. Fix $k \in \mathbb{N}$ for the rest of the paper. k will denote the tree-width bound we consider.

We introduce the notion of **oriented** k**-trees** which refines the notion of k-trees. Let $\mathcal{R} = \{R_1, \ldots, R_k\}$ be a vocabulary consisting of binary relation symbols. An oriented k-tree is an $\mathcal{R}$-structure $\mathfrak{R}$ in which all $R_i^{\mathfrak{R}}$ are total functions and whose Gaifman graph $\mathrm{Gaif}(\mathfrak{R})$ is a partial k-tree.

---

[2] On relational structures, MSO is also known as *Guarded Second Order logic* GSO. The results of this paper extend to CMSO, the extension of MSO with modular counting quantifiers.

[3] Since we explicitly allowed quantification over subsets of relations for MSO, we do not view graphs and structures as incidence structures, in contrast to [6, Sections 1.8.1 and 1.9.1].

Lemma 1. *Every $\mathcal{C}$-structure $\mathfrak{M}$ of tree-width k can be expanded into a $(\mathcal{C} \cup \mathcal{R})$-structure $\mathfrak{N}$ such that: (i) $\mathfrak{N}|_{\mathcal{R}}$ is an oriented k-tree, (ii) $\mathrm{Gaif}(\mathfrak{N})$ is a subgraph of $\mathrm{Gaif}(\mathfrak{N}|_{\mathcal{R}})$, and (iii) the tree-width of $\mathfrak{N}$ is k.*

The oriented 2-tree in Fig. 2(b) is an expansion of the directed graph in Fig. 2(a) as guaranteed in Lemma 1. In Fig. 2(b), R_1 and R_2 are denoted by the dashed arrows and the dotted arrows, respectively. There are several other oriented k-trees which expand Fig. 2(a) and fulfill the requirements in Lemma 1,

To see that Lemma 1 holds, we describe a construction of $\mathfrak{N}$ echoing the process of constructing k-trees above. For each vertex u of the initial $(k+1)$-clique, we can set the values of $R_1^{\mathfrak{N}}(u), \ldots, R_k^{\mathfrak{N}}(u)$ to be the other k vertices of the clique. When a new vertex u is added to the k-tree, k edges incident to it are added. We set $R_1^{\mathfrak{N}}(u), \ldots, R_k^{\mathfrak{N}}(u)$ to be the set of vertices incident to u. For oriented k-trees whose Gaifman graph is not a k-tree the construction of an oriented k-tree is augmented by changing the value of $R_i^{\mathfrak{N}}(u)$ to $R_i^{\mathfrak{N}}(u) = u$ whenever $R_i^{\mathfrak{N}}(u)$ is not well-defined. This can happen when the target of u under $R_i^{\mathfrak{N}}$ is a vertex which was eliminated by taking the subgraph of a k-tree to obtain the partial k-tree.

3. Overview of the Main Theorem and its Proof

The precise statement of the main theorem is as follows:

Theorem 1 (Main Theorem)**.** *Let $\mathcal{C}_{bnd}$ and $\mathcal{C}_{unb}$ be vocabularies. Let s be a binary relation symbol not in $\mathcal{C}_{bnd} \cup \mathcal{C}_{unb}$. Let $\alpha \in \mathrm{MSO}(\mathcal{C}_{bnd})$ and $\beta \in C^2(\mathcal{C}_{unb})$. There is an effectively computable sentence $\delta \in C^2(\mathcal{D})$ over a vocabulary $\mathcal{D} \supseteq \{s\}$ such that the following are equivalent:*

(i) There is a $(\mathcal{C}_{bnd} \cup \mathcal{C}_{unb})$-structure $\mathfrak{M}$ such that $\mathfrak{M} \models \alpha \wedge \beta$ and $\mathrm{tw}(\mathfrak{M}|_{\mathcal{C}_{bnd}}) \leq k$.
(ii) There is a $\mathcal{D}$-structure $\mathfrak{N}$ such that $\mathfrak{N} \models \delta$ and $s^{\mathfrak{N}}$ is a binary tree.

The first step towards proving Theorem 1 is the Separation Theorem:

Theorem 2 (Separation Theorem)**.** *Let $\mathcal{C}_{bnd}$ and $\mathcal{C}_{unb}$ be vocabularies. Let $\alpha \in \mathrm{MSO}(\mathcal{C}_{bnd})$ and $\beta \in C^2(\mathcal{C}_{unb})$. There are effectively computable sentences $\alpha' \in \mathrm{MSO}(\mathcal{D}_{bnd})$ and $\beta' \in C^2(\mathcal{D}_{unb})$ over vocabularies $\mathcal{D}_{bnd}$ and $\mathcal{D}_{unb}$ such that $\mathcal{D}_{bnd} \cap \mathcal{D}_{unb}$ only contains unary relation symbols and the following are equivalent:*

(i) There is a $(\mathcal{C}_{bnd} \cup \mathcal{C}_{unb})$-structure $\mathfrak{M}$ with $\mathfrak{M} \models \alpha \wedge \beta$ and $\mathrm{tw}(\mathfrak{M}|_{\mathcal{C}_{bnd}}) \leq k$
(ii) There is a $(\mathcal{D}_{bnd} \cup \mathcal{D}_{unb})$-structure $\mathfrak{N}$ with $\mathfrak{N} \models \alpha' \wedge \beta'$ and $\mathrm{tw}(\mathfrak{N}|_{\mathcal{D}_{bnd}}) \leq k$.

In conjunction with Theorem 2, we only need to prove Theorem 1 in the case that the MSO-formula α and the C^2-formula β only share unary relation symbols. The significance of Theorem 2 is that it allows us to use tools designed for MSO in our more involved setting. The proof of Theorem 2 uses notions of types for C^2-sentences in Scott normal form, oriented k-trees, coloring arguments, and an induction on ranks of structures. Theorem 2 is discussed in Section 4. The next step is to move from structures whose reducts have bounded tree-width to structures which contain a binary tree.

Lemma 2. *Let $\mathcal{C}_{bnd}$ and $\mathcal{C}_{unb}$ be vocabularies such that $\mathcal{C}_{bnd} \cap \mathcal{C}_{unb}$ contains only unary relation symbols. Let s be a binary relation symbol. There is a vocabulary $\mathcal{D}_{bnd}$ consisting of s and unary relation symbols only, and, for every $\alpha \in \mathrm{MSO}(\mathcal{C}_{bnd})$ and $\beta \in C^2(\mathcal{C}_{unb})$, effectively computable sentences $\alpha' \in \mathrm{MSO}(\mathcal{D}_{bnd})$ and $\beta' \in C^2(\mathcal{D}_{bnd} \cup \mathcal{C}_{unb})$ such that the following are equivalent:*

(i) There is a $(\mathcal{C}_{bnd} \cup \mathcal{C}_{unb})$*-structure* $\mathfrak{M}$ *such that* $\mathfrak{M} \models \alpha \wedge \beta$ *and such that* $\mathrm{tw}(\mathfrak{M}|_{\mathcal{C}_{bnd}}) \leq k$.

(ii) There is a $(\mathcal{D}_{bnd} \cup \mathcal{C}_{unb})$*-structure* $\mathfrak{N}$ *such that* $\mathfrak{N} \models \alpha' \wedge \beta'$ *and* $s^{\mathfrak{N}}$ *is a binary tree.*

Technically, Lemma 2 is proved using a translation scheme which maps structures with a binary tree into structures whose $\mathcal{C}_{bnd}$-reducts have tree-width at most k, and conversely, each of the latter structures is the image of a structure with a binary tree under the translation scheme. Translation schemes capturing the graphs of tree-width at most k as the image of labeled trees were studied in the context of decidability and model checking of MSO [1]. We need a more refined construction to ensure that the translation scheme also behaves correctly on C^2-sentences, i.e. that it maps C^2-sentences to C^2-sentences, see Lemma 2 in Section 5.

Now that we have reduced our attention to the case that our structures contain a binary tree, we can replace MSO-sentences with equi-satisfiable C^2-sentences.

Lemma 3. *Let* $\mathcal{C}$ *be a vocabulary which consists only of a binary relation symbol* s *and unary relation symbols. Let* α *be an* $\mathrm{MSO}(\mathcal{C})$*-sentence. There is an effectively computable* $C^2(\mathcal{D})$*-sentence* γ *over a vocabulary* $\mathcal{D} \supseteq \mathcal{C}$ *such that for every* $\mathcal{C}$*-structure* $\mathfrak{M}$ *in which* $s^{\mathfrak{M}}$ *is a binary tree the following are equivalent:*

(i) $\mathfrak{M} \models \alpha$ *(ii) There is a* $\mathcal{D}$*-structure* $\mathfrak{N}$ *expanding* $\mathfrak{M}$ *such that* $\mathfrak{N} \models \gamma$.

For the proof of Lemma 3 we use a Feferman-Vaught type theorem which states that the Hintikka type (i.e. MSO types) of a binary tree labeled with unary relation symbols depends only on the Hintikka types of its children. We can therefore axiomatize in C^2 that the Hintikka type of the labeled binary tree implies a given MSO-sentence.

Having replaced the MSO-sentence in statement (ii) of Lemma 2 with a C^2-sentence, we are left with the problem of deciding whether a C^2-sentence is satisfiable by a structure in which a specified relation is a binary tree, which has recently been shown to be decidable:

Theorem 3 (Charatonik and Witkowski [4]). *Let* $\mathcal{C}$ *be a vocabulary which contains a binary relation symbol* s*. Given a* $C^2(\mathcal{C})$*-sentence* φ*, it is decidable whether* φ *is satisfiable by a structure* $\mathfrak{M}$ *in which* $s^{\mathfrak{M}}$ *is a binary tree.*

4. Separation Theorem

4.1. Basic Definitions and Results.

1-types and 2-types. We begin with some notation and definitions in the spirit of the literature on decidability of C^2, cf. e.g. [22, 4]. Let $\mathcal{A}$ be a vocabulary of unary and binary relations.

A **1-type** π is a maximal consistent set of atomic $\mathcal{A}$-formulas or negations of atomic $\mathcal{A}$-formulas with free variable x, i.e., exactly one of $A(x)$ and $\neg A(x)$ belongs to π for every unary relation symbol $A \in \mathcal{A}$, and exactly one of $B(x,x)$ and $\neg B(x,x)$ belongs to π for every binary relation symbol $B \in \mathcal{A}$. We denote by $\mathfrak{char}_{\pi}(x) = \bigwedge_{\iota \in \pi} \iota$ the formula that characterizes the 1-type π. We denote by 1-Types($\mathcal{A}$) the set of 1-types over $\mathcal{A}$.

A **2-type** λ is a maximal consistent set of atomic $\mathcal{A}$-formulas or negations of atomic $\mathcal{A}$-formulas with free variables x and y, i.e., for every $z \in \{x, y\}$ and unary relation symbol $A \in \mathcal{A}$, exactly one of $A(z)$ and $\neg A(z)$ belongs to λ, and for every $z_1, z_2 \in \{x, y\}$ and binary relation symbol $B \in \mathcal{A}$, exactly one of $B(z_1, z_2)$ and $\neg B(z_1, z_2)$ belongs to λ. We note that the equality relation $\approx$ is also part of a 2-type. We write λ^{-1} for the 2-type obtained from λ by substituting all occurrences of x resp. y with y resp. x. We write λ_x for the 1-type obtained from λ by restricting λ to formulas with free variable x. We write λ_y for the 1-type obtained from λ by restricting λ to formulas with free variable y *and substituting* y *with* x. We denote by $\mathfrak{char}_\lambda(x, y) = \bigwedge_{\iota \in \lambda} \iota$ the formula that characterizes the 2-type λ. We denote by 2-Types($\mathcal{A}$) the set of 2-types over $\mathcal{A}$.

Let $\mathfrak{M}$ be a $\mathcal{A}$-structure. We denote by $1\text{-}\mathfrak{tp}^{\mathfrak{M}}(u)$ the unique 1-type π such that $\mathfrak{M} \models \mathfrak{char}_\pi(u)$. For elements u, v of $\mathfrak{M}$, we denote by $2\text{-}\mathfrak{tp}^{\mathfrak{M}}(u, v)$ the unique 2-type λ such that $\mathfrak{M} \models \mathfrak{char}_\lambda(u, v)$. We denote by $2\text{-}\mathfrak{tp}(\mathfrak{M}) = \{2\text{-}\mathfrak{tp}^{\mathfrak{M}}(u, v) \mid u, v \text{ elements of } \mathfrak{M}\}$ the set of 2-types *realized* by $\mathfrak{M}$. The following lemma is easy to see:

Lemma 4. *Let* $\mathfrak{M}_1, \mathfrak{M}_2$ *be two* $\mathcal{A}$*-structures over the same universe* M *and let* $\phi = \forall x, y.\, \chi \in C^2(\mathcal{A})$ *with* χ *quantifier-free. If* $2\text{-}\mathfrak{tp}(\mathfrak{M}_1) = 2\text{-}\mathfrak{tp}(\mathfrak{M}_2)$*, then* $\mathfrak{M}_1 \models \phi$ *iff* $\mathfrak{M}_2 \models \phi$*.*
(See proof in Appendix 8.4.)

Scott Normal Form and $\mathcal{T}$-functionality. C^2-sentences have a Scott-Normal Form, cf. [12], which can be obtained by iteratively applying Skolemization and introducing new predicates for subformulas, together with predicates ensuring the soundness of this transformation:

Lemma 5 (Scott Normal Form, [12])**.** *For every* C^2*-sentence* β *there is a* C^2*-sentence* β' *of the form*

$$\forall x, y.\, \chi \wedge \bigwedge_{i \in [l]} \forall x.\, \exists^{=1} y.\, S_i(x, y), \tag{1}$$

with χ *quantifier-free, over an expanded vocabulary such that* β *and* β' *are equi-satisfiable. Moreover,* β' *is computable. The expanded vocabulary contains in particular the fresh binary relation symbols in* $\mathcal{S} = \{S_1, \ldots, S_l\}$

Let $\mathcal{T}$ be a set of binary relation symbols. We say a structure $\mathfrak{M}$ is **$\mathcal{T}$-functional**, if for every $T \in \mathcal{T}$, $T^{\mathfrak{M}}$ is a total function on the universe of $\mathfrak{M}$. Observe the following are equivalent for every structure $\mathfrak{M}$:

(i) $\mathfrak{M}$ satisfies Eq. (1), and (ii) $\mathfrak{M} \models \forall x, y.\, \chi$ and $\mathfrak{M}$ is $\mathcal{S}$-functional.

Message Types and Chromaticity. Let $\mathcal{T} \subseteq \text{bin}(\mathcal{A})$ be a subset of the binary relation symbols of $\mathcal{A}$. We write $\lambda \in \mathcal{T}$-MsgTypes($\mathcal{A}$) and say λ is a **$\mathcal{T}$-message type**, if $\lambda \in$ 2-Types($\mathcal{A}$) and $T(x, y) \in \lambda$ for some $T \in \mathcal{T}$. Let $\mathfrak{M}$ be a $\mathcal{A}$-structure with universe M. We define $E^{\mathfrak{M}}_{\mathcal{T}} = \{(u, v) \in M^2 \mid \text{there is a } T \in \mathcal{T} \text{ with } \mathfrak{M} \models T(u, v)\}$. The **$\mathcal{T}$-message-graph** is the directed graph $G^{\mathfrak{M}}_{\mathcal{T}} = (M, E)$, where $E = \{(u, v) \in M^2 \mid u \neq v \text{ and } (u, v) \in E^{\mathfrak{M}}_{\mathcal{T}} \circ E^{\mathfrak{M}}_{\mathcal{T}}\}$, where $R \circ S = \{(a, b) \mid \text{there is a } c \text{ with } (a, c) \in R \text{ and } (c, b) \in S\}$ denotes the usual *composition* of relations. We say $\mathfrak{M}$ is **$\mathcal{T}$-chromatic**, if $1\text{-}\mathfrak{tp}^{\mathfrak{M}}(u) \neq 1\text{-}\mathfrak{tp}^{\mathfrak{M}}(v)$ for all $(u, v) \in E$.

We note that if $\mathfrak{M}$ is $\mathcal{T}$-functional, then $G^{\mathfrak{M}}_{\mathcal{T}}$ has out-degree $deg^+(u) \leq |\mathcal{T}|^2$ for all $u \in M$. This allows us to prove Lemma 6 based on Lemma 7; the proofs can be found in Appendices 8.6 and 8.5.

Lemma 6. *There is a finite set of unary relations symbols* $\mathrm{colors}_{\mathcal{T}}$ *such that every* $\mathcal{T}$*-functional* $\mathcal{A}$*-structure can be expanded to a* $\mathcal{T}$*-chromatic* $(\mathcal{A}\cup\mathrm{colors}_{\mathcal{T}})$*-structure.*

Lemma 7. *Let* $G=(V,E)$ *be a directed graph with out-degree* $deg^{+}(v)\leq k$ *for all* $v\in V$*. Then, the underlying undirected graph has a proper* $2k+1$*-coloring.*

4.2. **Separation Theorem.** Let $G=(V,E)$ be an (undirected) graph. We say G is k**-bounded**, if the edges of G can be oriented such that every node of G has out-degree less than k. We say a structure $\mathfrak{M}$ is k-bounded if its Gaifman graph is k-bounded.

Theorem 2 (Separation Theorem)**.** *Let* k *be a natural number. Let* $\mathcal{C}_{bnd}$ *and* $\mathcal{C}_{unb}$ *be vocabularies. Let* $\alpha\in\mathrm{MSO}(\mathcal{C}_{bnd})$ *and*[4] $\beta\in C^{2}(\mathcal{C}_{unb})$*. There are effectively computable sentences* $\alpha'\in\mathrm{MSO}(\mathcal{D}_{bnd})$ *and* $\beta'\in C^{2}(\mathcal{D}_{unb})$ *over vocabularies* $\mathcal{D}_{bnd}$ *and* $\mathcal{D}_{unb}$ *such that* $\mathcal{D}_{bnd}\cap\mathcal{D}_{unb}$ *only contains unary relation symbols such that for every* k*-bounded graph* G *the following are equivalent:*

(i) *There is a* $(\mathcal{C}_{bnd}\cup\mathcal{C}_{unb})$*-structure* $\mathfrak{M}$ *with* $\mathfrak{M}\models\alpha\wedge\beta$ *and* $\mathrm{Gaif}(\mathfrak{M}|_{\mathcal{C}_{bnd}})=G$*.*
(ii) *There is a* $(\mathcal{D}_{bnd}\cup\mathcal{D}_{unb})$*-structure* $\mathfrak{N}$ *with* $\mathfrak{N}\models\alpha'\wedge\beta'$ *and* $\mathrm{Gaif}(\mathfrak{N}|_{\mathcal{D}_{bnd}})=G$*.*

We assume that β is in the form given in Eq. (1) for some set of binary relation symbols $\mathcal{S}=\{S_1,\ldots,S_l\}\subseteq\mathcal{C}_{unb}$ and quantifier-free C^2-formula χ. Let $\mathcal{R}=\{R_1,\ldots,R_k\}$ be a set of fresh binary relation symbols. We set $\mathcal{T}=\mathcal{S}\cup\mathcal{R}$. We begin by giving an intuition for the proof of the Separation Theorem in three stages.

1. Syntactic separation coupled with semantic constraints. For a binary relation symbol B, we define its *copy* as the relation symbol $\overline{B}$. For every vocabulary $\mathcal{A}$, we define its *copy* $\overline{\mathcal{A}}=\mathrm{un}(\mathcal{A})\cup\{\overline{B}\mid B\in\mathcal{A}\}$ to be the unary relation symbols of $\mathcal{A}$ plus the copies of its binary relations symbols. We assume that copied relation symbols are distinct from non-copied symbols, i.e., $\mathrm{bin}(\mathcal{A})\cap\mathrm{bin}(\overline{\mathcal{A}})=\emptyset$. For a formula φ over vocabulary $\mathcal{A}$, we define its *copy* $\overline{\varphi}$ over vocabulary $\overline{\mathcal{A}}$ as the formula obtained from φ by substituting every occurrence of a binary relation symbol $B\in\mathcal{A}$ with $\overline{B}$.

The sentences $\overline{\alpha}$ (the copy of α) and β do not share any binary relation symbols. Clearly, *(i)* from Theorem 2 holds iff

(I) $\overline{\alpha}\wedge\beta$ is satisfied by a $((\overline{\mathcal{C}_{bnd}}\cup\overline{\mathcal{C}_{unb}})\cup(\mathcal{C}_{bnd}\cup\mathcal{C}_{unb}))$-structure $\mathfrak{N}$ with $B^{\mathfrak{N}}=\overline{B}^{\mathfrak{N}}$ for all $B\in\mathrm{bin}(\mathcal{C}_{bnd})$ and $\mathrm{Gaif}(\mathfrak{M}|_{\mathcal{C}_{bnd}})=G$.

In the next two stages we will construct α' and β' so that *(I)* is equivalent to *(ii)* from Theorem 2. More precisely, we will construct sentences $\mu_{bnd},\mu_{unb}\in C^2(\mathcal{D}_{unb})$ with $\mathcal{D}_{unb}\supseteq\mathcal{C}_{bnd}\cup\mathcal{C}_{unb}$ and $\mathcal{D}_{bnd}=\overline{\mathcal{D}_{unb}}$ such that *(I)* is equivalent *(II)*:

(II) $(\overline{\alpha}\wedge\overline{\mu_{bnd}})\wedge(\beta\wedge\mu_{unb})$ is satisfied by a $(\mathcal{D}_{bnd}\cup\mathcal{D}_{unb})$-structure $\mathfrak{N}$ with $\mathrm{Gaif}(\mathfrak{N}|_{\mathcal{D}_{bnd}})=G$.

2. Representation of k**-bounded structures using functions and unary relations.** Theorem 2 as well as *(I)* and *(II)* involve reducts which are k-bounded structures. k-bounded $\mathcal{A}$-structures $\mathfrak{A}$ can be represented by introducing new binary relation symbols interpreted as functions and new unary relation symbols as follows.

(a) We add k fresh relation symbols $\mathcal{R}=\{R_1,\ldots,R_k\}$ and axiomatize that these relations are interpreted as total functions.

---

[4] The Separation Theorem remains correct if we replace C^2 with any logic containing C^2 which is closed under conjunction.

(b) We add fresh unary relations $\{P_\lambda \mid \lambda \in \mathcal{R}\text{-MsgTypes}(\mathcal{A})\}$ and axiomatize that every element labeled by P_λ has an outgoing edge with 2-type λ. The symbols P_λ are called *unary 2-type annotations*.

(c) We axiomatize that $\mathrm{Gaif}(\mathfrak{A}|_{\mathrm{bin}(\mathcal{A})}) = \mathrm{Gaif}(\mathfrak{A}|_{\mathcal{R}})$.

In other words, the functions interpreting $R_1, \ldots, R_k$ witness that $\mathfrak{A}$ can be oriented so that every node in the Gaifman graph of $\mathfrak{A}$ has outdegree at most k. The 2-type of each edge (u, v) in $\mathfrak{A}$ is encode by putting the unary relation symbol P_λ of the 2-type of (u, v) on the source u in the orientation.

Given a $((\overline{\mathcal{C}_{bnd}} \cup \overline{\mathcal{C}_{unb}}) \cup (\mathcal{C}_{bnd} \cup \mathcal{C}_{unb}))$-structure $\mathfrak{N}$, we will use the above representation twice, on $\mathfrak{N}|_{\mathcal{C}_{bnd}}$ and $\mathfrak{N}|_{\overline{\mathcal{C}_{bnd}}}$, by axiomatizing that every element labeled by P_λ has an outgoing edge with 2-type λ and an outgoing edges with 2-type $\overline{\lambda}$. This will allow us to replace the condition from *(I)* that $B^{\mathfrak{N}} = \overline{B}^{\mathfrak{N}}$ for all $B \in \mathrm{bin}(\mathcal{C}_{bnd})$ with the condition that $R_i^{\mathfrak{N}} = \overline{R_i}^{\mathfrak{N}}$ for all $R_i \in \mathcal{R}$. We will define a vocabulary $\mathcal{E} \supseteq \mathcal{C}_{bnd} \cup \mathcal{C}_{unb} \cup \mathcal{R}$. Let $\mathcal{P}_{\mathcal{R}} = \{P_\lambda \mid \lambda \in \mathcal{R}\text{-MsgTypes}(\mathcal{E})\}$. We will define two vocabularies $\mathcal{D}_{unb} \supseteq \mathcal{C}_{bnd} \cup \mathcal{C}_{unb} \cup \mathcal{R} \cup \mathcal{P}_{\mathcal{R}}$ and $\mathcal{D}_{bnd} = \overline{\mathcal{D}_{unb}}$. According to (a), (b), and (c), we will construct $\nu_{bnd}, \nu_{unb} \in C^2(\mathcal{D}_{unb})$ such that *(I)* is equivalent to the following:

(I') $(\overline{\alpha} \wedge \overline{\nu_{bnd}}) \wedge (\beta \wedge \nu_{unb})$ is satisfied by a $\mathcal{D}_{bnd} \cup \mathcal{D}_{unb}$-structure $\mathfrak{M}$ with $R_i^{\mathfrak{M}} = \overline{R_i}^{\mathfrak{M}}$ for all $R_i \in \mathcal{R}$ and $\mathrm{Gaif}(\mathfrak{M}|_{\mathcal{R}}) = G$.

3. Establishing the semantic condition of *(I')* by swapping edges. Here we discuss how to show the implication from *(II)* to *(I')* and make the vocabularies $\mathcal{D}_{bnd}$ and $\mathcal{D}_{unb}$ precise. Let $\mathfrak{N}$ be a $(\mathcal{D}_{bnd} \cup \mathcal{D}_{unb})$-structure with $\mathfrak{N} \models (\overline{\alpha} \wedge \overline{\mu_{bnd}}) \wedge (\beta \wedge \mu_{unb})$. It simplifies the discussion to split a $(\mathcal{D}_{bnd} \cup \mathcal{D}_{unb})$-structure $\mathfrak{N}$ into two $\mathcal{D}_{unb}$-structures. The $\mathcal{D}_{unb}$-structure $\mathfrak{L}$ is $\mathfrak{N}|_{\mathcal{D}_{unb}}$. The $\mathcal{D}_{unb}$-structure $\mathfrak{L}'$ is obtained from $\mathfrak{N}|_{\mathcal{D}_{bnd}}$ by renaming copies of relation symbols $\overline{B}$ to B — i.e., we define the $\mathcal{D}_{unb}$-structure $\mathfrak{L}'$ by setting $1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{N}}(u)$ for all $u \in M$ and setting $\mathfrak{L}' \models B(u, v)$ iff $\mathfrak{N} \models \overline{B}(u, v)$ for all $u, v \in M$ and $B \in \mathrm{bin}(\mathcal{D}_{bnd})$. The interpretations of the relations R_i might differ in $\mathfrak{L}$ and $\mathfrak{L}'$. Observe that we have $\mathfrak{L}' \models \alpha$ and $\mathfrak{L} \models \beta$. The *key idea of the proof* is prove the existence of a sequence of structures $\mathfrak{L} = \mathfrak{L}_0, \ldots, \mathfrak{L}_p$, where each $\mathfrak{N}_{i+1}$ is obtained from $\mathfrak{N}_i$ by *swapping edges*, until the interpretations of the relations R_i agree in $\mathfrak{L}_p$ and $\mathfrak{L}'$. The edge swapping operation is a local operation which involves changing the 2-types of at most 4 edges.

The edge swapping operation satisfies two crucial preservation requirements: edge swapping preserves (PR-1) the truth value of β, i.e. $\mathfrak{L}_p \models \beta$, and (PR-2) $\mathcal{R}$-functionality. The universal constraint $\forall x, y.\, \chi$ in β is maintained under edge swapping because this operation does not change the set of 2-types (see Lemma 4). To satisfy the preservation requirements (PR-1) and (PR-2), all that remains is to guarantee the existence of a sequence of edge swapping preserving $\mathcal{T}$-functionality. Note that $\mathcal{T}$-functionality amounts to $\mathcal{R}$-functionality and $\mathcal{S}$-functionality. We use two main techqniues for ensuring the preservation of $\mathcal{T}$-functionality: chromaticity and unary 2-type annotations. We will axiomatize that the structures $\mathfrak{L}$ and $\mathfrak{L}'$ are chromatic and we will take care that chromaticity is maintained during edge swaps. We will add fresh unary relation symbols $\mathcal{P}_{\mathcal{S}} = \{P_\lambda \mid \lambda \in \mathcal{S}\text{-MsgTypes}(\mathcal{E})\}$ and axiomatize that every element of $\mathfrak{N}$ labeled by P_λ has an outgoing edge with 2-type λ and an outgoing edge with 2-type $\overline{\lambda}$.

Proof of the Separation Theorem. We now start the formal proof of the Separation Theorem. Let $\mathrm{colors}_{\mathcal{T}}$ be the vocabulary from Lemma 6. We set $\mathcal{E} = \mathcal{C}_{bnd} \cup \mathcal{C}_{unb} \cup \mathcal{R} \cup \mathrm{colors}_{\mathcal{T}}$. We set $\mathcal{D}_{bnd} = \mathcal{E} \cup \mathcal{P}_{\mathcal{S}} \cup \mathcal{P}_{\mathcal{R}}$ and $\mathcal{D}_{unb} = \overline{\mathcal{D}_{bnd}}$. Next we will define formulas $\psi \in \mathrm{MSO}(\mathcal{D}_{bnd})$ and $\beta' \in C^2(\mathcal{D}_{bnd})$, and set $\alpha' = \overline{\psi} \in \mathrm{MSO}(\mathcal{D}_{unb})$. We set $\nu_{bnd} = \delta \wedge \theta \wedge \epsilon \wedge \zeta_{bnd}$, $\nu_{unb} = \delta \wedge \theta \wedge \zeta_{unb}$, $\mu_{bnd} = \nu_{bnd} \wedge \eta$, and $\mu_{unb} = \nu_{unb} \wedge \eta$, $\psi = \alpha \wedge \mu_{bnd}$, and $\beta' = \beta \wedge \mu_{unb}$, where:

$$
\begin{aligned}
\delta &= \textstyle\bigwedge_{i \in [k]} \forall x.\, \exists^{=1} y.\, R_i(x,y) \\
\theta &= \forall x, y.\, x \not\approx y \rightarrow \Big(\textstyle\bigvee_{i \in [k]} R_i(x,y) \vee R_i(y,x) \leftrightarrow \bigvee_{B \in \mathrm{bin}(\mathcal{C}_{bnd})} B(x,y) \vee B(y,x) \Big) \\
\epsilon &= \textstyle\bigwedge_{B \in \mathrm{bin}(\mathcal{C}_{unb})} \forall x, y.\, x \not\approx y \rightarrow \Big(B(x,y) \rightarrow \bigvee_{i \in [k]} R_i(x,y) \vee R_i(y,x) \Big)
\end{aligned}
$$

$$
\zeta_{bnd} = \bigwedge_{\substack{\lambda \in \mathcal{R}\text{-MsgTypes}(\mathcal{E}) \text{ or} \\ (\lambda \in \mathcal{T}\text{-MsgTypes}(\mathcal{E}) \text{ and } \lambda^{-1} \in \mathcal{R}\text{-MsgTypes}(\mathcal{E}))}} \forall x.\, P_\lambda(x) \leftrightarrow \exists y. \mathfrak{char}_\lambda(x,y)
$$

$$
\zeta_{unb} = \bigwedge_{\lambda \in \mathcal{T}\text{-MsgTypes}(\mathcal{E})} \forall x.\, P_\lambda(x) \leftrightarrow \exists y. \mathfrak{char}_\lambda(x,y)
$$

$$
\eta = \bigwedge_{\substack{\lambda \in \mathcal{T}\text{-MsgTypes}(\mathcal{E}) \\ T, T' \in \mathcal{T} \,:\, T'(x,y) \in \lambda}} \forall x, y.\, P_\lambda(x) \wedge T(y,x) \wedge \neg T'(x,y) \rightarrow \neg \mathfrak{char}_{\lambda_y}(y)
$$

δ expresses that each R_i is interpreted as a total function, θ expresses that for every $\mathcal{D}_{unb}$-structure $\mathfrak{L}$ with $\mathfrak{L} \models \theta$ we have that $\mathrm{Gaif}(\mathfrak{L}|_{\mathcal{R}}) = \mathrm{Gaif}(\mathfrak{L}|_{\mathcal{C}_{bnd}})$, ϵ expresses that for every $\mathcal{D}_{unb}$-structure $\mathfrak{L}$ with $\mathfrak{L} \models \epsilon$ we have that $\mathrm{Gaif}(\mathfrak{L}|_{\mathcal{D}_{unb}})$ is a subgraph of $\mathrm{Gaif}(\mathfrak{L}|_{\mathcal{R}})$, ζ expresses that for every $\lambda \in \mathcal{T}\text{-MsgTypes}(\mathcal{E})$ with $\lambda \in \mathcal{R}\text{-MsgTypes}(\mathcal{E})$ or $\lambda \in \mathcal{R}\text{-MsgTypes}(\mathcal{E})$, $\mathcal{D}_{unb}$-structure $\mathfrak{L}$ and element u of $\mathfrak{L}$ we have $\mathfrak{L} \models P_\lambda(u)$ iff there is an element v of $\mathfrak{L}$ such that $2\text{-}\mathfrak{tp}^{\mathfrak{L}|_{\mathcal{E}}}(u,v) = \lambda$, ζ expresses that for every $\lambda \in \mathcal{T}\text{-MsgTypes}(\mathcal{E})$, $\mathcal{D}_{unb}$-structure $\mathfrak{L}$ and element u of $\mathfrak{L}$ we have $\mathfrak{L} \models P_\lambda(u)$ iff there is an element v of $\mathfrak{L}$ such that $2\text{-}\mathfrak{tp}^{\mathfrak{L}|_{\mathcal{E}}}(u,v) = \lambda$, η expresses that for every $\mathcal{D}_{unb}$-structure $\mathfrak{L}$ with $\mathfrak{L} \models \zeta_{unb}$, $\mathfrak{L}$ is chromatic iff $\mathfrak{L} \models \eta$.

The direction "(i) implies (ii)" of the Separation Theorem is not hard:

Lemma 8. *Let G be a k-bounded graph. Let $\mathfrak{M}$ be a $(\mathcal{C}_{bnd} \cup \mathcal{C}_{unb})$-structure with $\mathfrak{M} \models \alpha \wedge \beta$ and $\mathrm{Gaif}(\mathfrak{M}|_{\mathcal{C}_{bnd}}) = G$. $\mathfrak{M}$ can be expanded to a $(\mathcal{D}_{bnd} \cup \mathcal{D}_{unb})$-structure $\mathfrak{N}$ with $\mathfrak{N} \models \alpha' \wedge \beta'$ and $\mathrm{Gaif}(\mathfrak{N}|_{\mathcal{D}_{bnd}}) = G$.*

Proof. Because of $\mathrm{Gaif}(\mathfrak{M}|_{\mathcal{C}_{bnd}}) = G$ and G is k-bounded, we can expand $\mathfrak{M}$ to a $\mathcal{C}_{bnd} \cup \mathcal{C}_{unb} \cup \mathcal{R}$-structure $\widehat{\mathfrak{M}}$ such that $\mathrm{Gaif}(\mathfrak{L}|_{\mathcal{R}}) = \mathrm{Gaif}(\mathfrak{L}|_{\mathcal{C}_{bnd}})$ and $R_i^{\widehat{\mathfrak{M}}}$ is a total function for all $i \in [k]$ (possibly adding self-loops for the relations R_i). Thus, $\widehat{\mathfrak{M}} \models \delta \wedge \theta$. According to Lemma 6, $\widehat{\mathfrak{M}}$ can be expanded to a chromatic structure $\widetilde{\mathfrak{M}}$ over vocabulary $\mathcal{E}$ with $\widetilde{\mathfrak{M}} \models \eta$. We expand $\widetilde{\mathfrak{M}}$ to a $\mathcal{D}_{bnd}$-structure $\mathfrak{L}$ such that for all $u \in M$ and $\lambda \in \mathcal{T}\text{-MsgTypes}(\mathcal{E})$ we have $\mathfrak{L} \models P_\lambda(u)$ iff there is an element v of $\mathfrak{L}$ such that $2\text{-}\mathfrak{tp}^{\mathfrak{L}|_{\mathcal{E}}}(u,v) = \lambda$. This definition gives us $\mathfrak{L} \models \zeta_{unb}$, and thus $\mathfrak{L} \models \beta'$. We expand $\mathfrak{L}$ to a $(\mathcal{D}_{bnd} \cup \mathcal{D}_{unb})$-structure $\mathfrak{N}$ such that for all $u, v \in M$ and $B \in \mathcal{B}$ we have $\mathfrak{N} \models \overline{B}(u,v)$ iff $\mathfrak{N} \models B(u,v)$ and $\mathfrak{N} \models R_i(u,v)$ or $\mathfrak{N} \models R_i(v,u)$ for some $i \in [k]$. We note that $\mathfrak{N} \models \alpha'$. ☐

Now we turn to the direction "(ii) implies (i)". Let G be a k-bounded graph. Let $\mathfrak{N}$ be a $(\mathcal{D}_{bnd} \cup \mathcal{D}_{unb})$-structure with $\mathfrak{N} \models \alpha' \wedge \beta'$ and $\mathrm{Gaif}(\mathfrak{N}|_{\mathcal{D}_{bnd}}) = G$. Let M be the universe of $\mathfrak{N}$. We define the $\mathcal{D}_{unb}$-structure $\mathfrak{L}'$ by setting $1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{N}}(u)$ for all $u \in M$ and setting $\mathfrak{L}' \models B(u,v)$ iff $\mathfrak{N} \models \overline{B}(u,v)$ for all $u,v \in M$ and $B \in \mathrm{bin}(\mathcal{D}_{bnd})$. We note that $\mathfrak{L}' \models \psi$ and $\mathrm{Gaif}(\mathfrak{L}') = G$. We define the $\mathcal{D}_{unb}$-structure $\mathfrak{L}$ by setting $\mathfrak{L} = \mathfrak{N}|_{\mathcal{D}_{unb}}$. We note that $\mathfrak{L} \models \beta'$.

We make the following definition: For $u \in M$ and $i \in [k]$ we set $rank^i_u(\mathfrak{L}, \mathfrak{L}') = 1$, if there are $v, w \in M$ with $\mathfrak{L} \models R_i(u,v)$, $\mathfrak{L}' \models R_i(u,w)$ and $v \neq w$; we set $rank^i_u(\mathfrak{L}, \mathfrak{L}') = 0$, otherwise. We set $rank_u(\mathfrak{L}, \mathfrak{L}') = \sum_{i \in [k]} rank^i_u(\mathfrak{L}, \mathfrak{L}')$ and $rank(\mathfrak{L}, \mathfrak{L}') = \sum_{u \in M} rank_u(\mathfrak{L}, \mathfrak{L}')$. *rank* measures the deviation of the relations $\mathcal{R}$ in $\mathfrak{L}$ and $\mathfrak{L}'$ (we note that there always are unique $v, w \in M$ for $u \in M$ with $\mathfrak{L} \models R_i(u,v)$, $\mathfrak{L}' \models R_i(u,w)$ because of $\mathfrak{L} \models \delta$ and $\mathfrak{L}' \models \delta$) and has the following important property:

Lemma 9. *If $rank(\mathfrak{L}, \mathfrak{L}') = 0$, then $\mathfrak{L}|_{\mathcal{C}_{bnd} \cup \mathcal{C}_{unb}} \models \alpha \wedge \beta$ and $\mathrm{Gaif}(\mathfrak{L}_{\mathcal{C}_{bnd}}) = G$.*
(See Appendix 8.8 for the proof.)

The proof of Lemma 9 uses the following simple but useful property of the rank function:

Lemma 10. *Let $u \in M$ be an element with $rank^i_u(\mathfrak{L}, \mathfrak{L}') = 0$ and let $\lambda \in$ 2-Types$(\mathcal{E})$ with $R_i \in \lambda$. For all $v \in M$ we have $2\text{-}\mathfrak{tp}^{\mathfrak{L}|_{\mathcal{E}}}(u,v) = \lambda$ iff $2\text{-}\mathfrak{tp}^{\mathfrak{L}'|_{\mathcal{E}}}(u,v) = \lambda$.*
(See Appendix 8.7 for the proof.)

Lemma 11. *There is a sequence of $\mathcal{D}_{unb}$-structures $\mathfrak{L}_0, \ldots, \mathfrak{L}_p$, with universe M and $\mathfrak{L} = \mathfrak{L}_0$, such that:*
(1) $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(u)$ for all $u \in M$,
(2) $2\text{-}\mathfrak{tp}(\mathfrak{L}_i) = 2\text{-}\mathfrak{tp}(\mathfrak{L}_{i+1})$ for all $0 \leq i < p$,
(3) $\mathfrak{L}_i \models \psi$, (in particular $\mathfrak{L}_i$ is $\mathcal{T}$-functional and chromatic),
(4) $rank(\mathfrak{L}_i, \mathfrak{L}') > rank(\mathfrak{L}_{i+1}, \mathfrak{L}')$ for all $0 \leq i < p$, and $rank(\mathfrak{L}_p, \mathfrak{L}') = 0$.

Proof. Assume we have already defined $\mathfrak{L}_i$ and $rank(\mathfrak{L}_i, \mathfrak{L}') > 0$. In the following we will define $\mathfrak{L}_{i+1}$. Because of $rank(\mathfrak{L}_i, \mathfrak{L}') > 0$ we can choose some elements $u, v, w \in M$ and $j \in [k]$ such that $\mathfrak{L}_i \models R_j(u,v)$, $\mathfrak{L}' \models R_j(u,w)$ and $v \neq w$. Let $\lambda = 2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u,v)$. We have $\lambda \in \mathcal{T}\text{-MsgTypes}(\mathcal{E})$ because of $R_j \in \mathcal{R}$. We have $\mathfrak{L}_i \models P_\lambda(u)$ because of $\mathfrak{L}_i \models \zeta_{unb}$. Because $1\text{-}\mathfrak{tp}(u)^{\mathfrak{L}_i} = 1\text{-}\mathfrak{tp}(u)^{\mathfrak{L}'}$ we have $\mathfrak{L}' \models P_\lambda(u)$. Thus, $\mathfrak{L}_i \models \mathfrak{char}_{\lambda_y}(v)$ and $\mathfrak{L}' \models \mathfrak{char}_{\lambda_y}(w)$. With $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i}(w) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(w)$ we get $\mathfrak{L}_i \models \mathfrak{char}_{\lambda_y}(w)$, and thus $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(v) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(w)$. We proceed by a case distinction:

Case 1: λ^{-1} is a $\mathcal{T}$-message type.

We have $\mathfrak{L}' \models P_{\lambda^{-1}}(w)$ because of $\mathfrak{L}' \models \zeta_{bnd}$. We get $\mathfrak{L}_i \models P_{\lambda^{-1}}(w)$ because of $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i}(w) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(w)$. With $\mathfrak{L}_i \models \zeta_{unb}$, there is an element $a \in M$ such that $\lambda = 2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(a,w)$ and $\mathfrak{L}_i \models P_\lambda(a)$. We note that $u \neq a$ because of $\mathfrak{L}_i \models R_j(a,w)$, $v \neq w$ and v is the unique element with $\mathfrak{L}_i \models R_j(u,v)$ (using $\mathfrak{L}_i \models \delta$). Moreover, $\mathfrak{L}_i \models \mathfrak{char}_{\lambda_y^{-1}}(a)$ and $\mathfrak{L}' \models \mathfrak{char}_{\lambda_y^{-1}}(u)$. With $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(u)$ we get $\mathfrak{L}_i \models \mathfrak{char}_{\lambda_y^{-1}}(u)$, and thus $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(a)$.

We note that the edges (u,w), (w,u), (a,v) and (v,a) do not have $\mathcal{T}$-message types (*) because $\mathfrak{L}_i$ is chromatic, $u \neq a$, $v \neq w$, $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(v) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(w)$, $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(a)$ and the edges (u,v), (v,u), (a,w) and (w,a) have $\mathcal{T}$-message types. Similarly, we get $\lambda \neq 2\text{-}\mathfrak{tp}^{\mathfrak{L}'|_{\mathcal{E}}}(u,v)$ and $\lambda \neq 2\text{-}\mathfrak{tp}^{\mathfrak{L}'|_{\mathcal{E}}}(a,w)$ (**) because $\mathfrak{L}'$ is chromatic, $\lambda = 2\text{-}\mathfrak{tp}^{\mathfrak{L}'|_{\mathcal{E}}}(v,w)$ and λ and λ^{-1} are $\mathcal{T}$-message types.

We define $\mathfrak{L}_{i+1}$ as follows: The unary relations of $\mathfrak{L}_{i+1}$ are defined such that we have $1\text{-}\mathfrak{tp}(u)^{\mathfrak{L}_{i+1}} = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i}(u)$ for all elements $u \in M$. We obtain the binary relations of $\mathfrak{L}_{i+1}$ by swapping the edges (u,w) and (u,v) as well as (a,w) and (a,v) in $\mathfrak{L}_i$: we set $2\text{-}\mathfrak{tp}^{\mathfrak{L}_{i+1}|_{\mathcal{E}}}(u,w) = \lambda$, $2\text{-}\mathfrak{tp}^{\mathfrak{L}_{i+1}|_{\mathcal{E}}}(a,v) = \lambda$, $2\text{-}\mathfrak{tp}^{\mathfrak{L}_{i+1}|_{\mathcal{E}}}(a,w) = 2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(a,v)$ and $2\text{-}\mathfrak{tp}^{\mathfrak{L}_{i+1}|_{\mathcal{E}}}(u,v) = 2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u,w)$; these 2-types are well-defined because of $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(v) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(w)$. All other 2-types in $\mathfrak{L}_{i+1}|_{\mathcal{E}}$ are the same as in $\mathfrak{L}_i|_{\mathcal{E}}$. This completes the definition of $\mathfrak{L}_{i+1}$.

We now argue that $\mathfrak{L}_{i+1}$ satisfies properties (1)-(4). Clearly, $\mathfrak{L}_{i+1}$ satisfies (1) by definition. Because we only swapped edges from $\mathfrak{L}_i$ to $\mathfrak{L}_{i+1}$ we have (2). From (2), Lemma 4 and $\mathfrak{L}_i \models \chi$, we get $\mathfrak{L}_{i+1} \models \chi$. Because we only swapped swapped edges from $\mathfrak{L}_i$ to $\mathfrak{L}_{i+1}$ we get $\mathfrak{L}_{i+1} \models \theta$ and $\mathfrak{L}_{i+1} \models \epsilon$ from $\mathfrak{L}_i \models \theta$ and $\mathfrak{L}_i \models \epsilon$. Because of (*) and $2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u,v) = \lambda = 2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(a,w)$ we get $\mathfrak{L}_{i+1} \models \zeta_{unb}$ from $\mathfrak{L}_i \models \zeta_{unb}$, that $\mathfrak{L}_{i+1}$ is $\mathcal{T}$-functional because $\mathfrak{L}_i$ is $\mathcal{T}$-functional and $\mathfrak{L}_{i+1} \models \eta$ from $\mathfrak{L}_i \models \eta$ (using that $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(v) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(w)$ and $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(a)$). Thus, $\mathfrak{L}_{i+1}$ satisfies property (3).

It remains to show property (4). Using (*), (**) and Lemma 10 we get that $rank_u(\mathfrak{L}_{i+1}, \mathfrak{L}') < rank_u(\mathfrak{L}_i, \mathfrak{L}')$ and $rank_z(\mathfrak{L}_{i+1}, \mathfrak{L}') \leq rank_z(\mathfrak{L}_i, \mathfrak{L}')$ for $z \in \{v, w, a\}$. Moreover, $rank_z(\mathfrak{L}_{i+1}, \mathfrak{L}') = rank_z(\mathfrak{L}_i, \mathfrak{L}')$ for $z \in M \setminus \{u, v, w, a\}$. Thus, we have $rank(\mathfrak{L}_i, \mathfrak{L}') > rank(\mathfrak{L}_{i+1}, \mathfrak{L}')$.

Case 2: λ^{-1} is not a $\mathcal{T}$-message type.

We note that the edge (w,u) does not have a $\mathcal{T}$-message type because $\mathfrak{L}_i$ is chromatic, $v \neq w$, $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(v) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(w)$ and the edge (u,v) has a $\mathcal{T}$-message type.

We define $\mathfrak{L}_{i+1}$ as follows: The unary relations of $\mathfrak{L}_{i+1}$ are defined such that we have $1\text{-}\mathfrak{tp}(u)^{\mathfrak{L}_{i+1}} = 1\text{-}\mathfrak{tp}(u)^{\mathfrak{L}_i}$ for all elements $u \in M$. We obtain the binary relations of $\mathfrak{L}_{i+1}$ by swapping edges in $\mathfrak{L}_i$: $2\text{-}\mathfrak{tp}^{\mathfrak{L}_{i+1}|_{\mathcal{E}}}(u,w) = \lambda$, and $2\text{-}\mathfrak{tp}^{\mathfrak{L}_{i+1}|_{\mathcal{E}}}(u,v) = 2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u,w)$. These 2-types are well-defined because of $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(v) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(w)$. All other 2-types in $\mathfrak{L}_{i+1}|_{\mathcal{E}}$ are the same as in $\mathfrak{L}_i|_{\mathcal{E}}$. This completes the definition of $\mathfrak{L}_{i+1}$.

We now argue that $\mathfrak{L}_{i+1}$ satisfies properties (1)-(4). As in the previous case, one can argue that $\mathfrak{L}_{i+1}$ satisfies (1) and (2), $\mathfrak{L}_{i+1} \models \chi$, $\mathfrak{L}_{i+1} \models \theta$ and $\mathfrak{L}_{i+1} \models \epsilon$. Because (v,u) and (w,u) do not have $\mathcal{T}$-message types we get $\mathfrak{L}_{i+1} \models \zeta_{unb}$ from $\mathfrak{L}_i \models \zeta_{unb}$ and that $\mathfrak{L}_{i+1}$ is $\mathcal{T}$-functional because $\mathfrak{L}_i$ is $\mathcal{T}$-functional. We get $\mathfrak{L}_{i+1} \models \eta$ from $\mathfrak{L}_i \models \eta$, $\mathfrak{L}' \models \eta$ and $1\text{-}\mathfrak{tp}^{\mathfrak{L}_i}(w) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(w)$.

It remains to show property (4). From Lemma 10 we get $2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(u,w) \neq \lambda$ and $2\text{-}\mathfrak{tp}^{\mathfrak{L}'|_{\mathcal{E}}}(u,v) \neq \lambda$. Again applying Lemma 10 we get $rank_u(\mathfrak{L}_{i+1}, \mathfrak{L}') < rank_u(\mathfrak{L}_i, \mathfrak{L}')$. Because $2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(v,u)$ and $2\text{-}\mathfrak{tp}^{\mathfrak{L}_i|_{\mathcal{E}}}(w,u)$ are not $\mathcal{T}$-message types, we get $rank_v(\mathfrak{L}_{i+1}, \mathfrak{L}') = rank_v(\mathfrak{L}_i, \mathfrak{L}')$ and $rank_w(\mathfrak{L}_{i+1}, \mathfrak{L}') = rank_w(\mathfrak{L}_i, \mathfrak{L}')$. Moreover, we have that $rank_z(\mathfrak{L}_{i+1}, \mathfrak{L}') = rank_z(\mathfrak{L}_i, \mathfrak{L}')$ for all $z \in M \setminus \{u, v, w\}$. Thus, $rank(\mathfrak{L}_i, \mathfrak{L}') > rank(\mathfrak{L}_{i+1}, \mathfrak{L}')$.

$\square$

5. FROM BOUNDED TREE-WIDTH TO BINARY TREES

This section is devoted to a discussion of the proof of Lemma 2. First we need some background from the literature. A **translation scheme for $\mathcal{C}_2$ over $\mathcal{C}_1$** is a tuple $\mathbf{t} = \langle \phi, \psi_C : C \in \mathcal{C}_2 \rangle$ of MSO($\mathcal{C}_1$)-formulas such that ϕ has exactly one free first order variable and the number of free first order variables in each

ψ_C is $arity(C)$. The formulas ϕ and ψ_C, $C \in \mathcal{C}_2$, do not have any free second order variables.[5] The **quantifier rank** $\mathrm{qr}(\mathbf{t})$ **of** $\mathbf{t}$ is the maximum of the quantifier ranks of ϕ and the ψ_C. $\mathbf{t}$ is **quantifier-free** if $\mathrm{qr}(\mathbf{t}) = 0$. The **induced transduction** $\mathbf{t}^\star$ is a partial function from $\mathcal{C}_1$-structures to $\mathcal{C}_2$-structures which assigns a $\mathcal{C}_2$-structure $\mathbf{t}^\star(\mathfrak{A})$ to a $\mathcal{C}_1$-structure $\mathfrak{A}$ as follows. The universe of $\mathbf{t}^\star(\mathfrak{A})$ is $A_\mathbf{t} = \{a \in A : \mathfrak{A} \models \phi(a)\}$. The interpretation of $C \in \mathcal{C}_2$ in $\mathbf{t}^\star(\mathfrak{A})$ is $C^{\mathbf{t}^\star(\mathfrak{A})} = \left\{\bar{a} \in {A_\mathbf{t}}^{arity(C)} : \mathfrak{A} \models \psi_C(\bar{a})\right\}$. Due to the convention that structures do not have an empty universe, $\mathbf{t}^\star(\mathfrak{A})$ is defined iff $\mathfrak{A} \models \exists x \phi(x)$.

Lemma 12 (Fundamental property of translation schemes)**.** *Let* $\mathbf{t}$ *be a translation scheme for* $\mathcal{C}_2$ *over* $\mathcal{C}_1$*. There is a computable function* $\mathbf{t}^\sharp$ *from* $\mathrm{MSO}(\mathcal{C}_2)$*-sentences to* $\mathrm{MSO}(\mathcal{C}_1)$*-sentences such that for every* $\mathcal{C}_1$*-structure* $\mathfrak{A}$ *for which* $\mathbf{t}^\star(\mathfrak{A})$ *is defined and for every* $\mathrm{MSO}(\mathcal{C}_2)$*-sentence* θ*,* $\mathfrak{A} \models \mathbf{t}^\sharp(\theta)$ *if and only if* $\mathbf{t}^\star(\mathfrak{A}) \models \theta$*.* $\mathbf{t}^\sharp$ *is called the* **induced translation**.

For an $\mathrm{MSO}(\mathcal{C}_2)$-sentence ζ, $\mathbf{t}^\sharp$ substitutes the relation symbols $C \in \mathcal{C}_2$ in ζ with the formulas ψ_C, requires that each of the free variables satisfies ϕ, and relativizes the quantification to ϕ. Appendix 8.9 gives an inductive definition of $\mathbf{t}^\sharp$ following Definition 3.2 of [19].

From the definition of the induced translation we have:

Lemma 13 (Quantifier-free translation schemes and C^2)**.** *Let* $\mathbf{t}$ *be a quantifier-free translation scheme for* $\mathcal{C}_2$ *over* $\mathcal{C}_1$*. The induced translation* $\mathbf{t}^\sharp$ *maps* $C^2(\mathcal{C}_2)$*-formulas to* $C^2(\mathcal{C}_1)$*-formulas.*

It is well-known that the class of graphs of tree-width at most k is the image of an induced transduction on a class of labeled trees [1]. For the proof of Lemma 2 we need an analogous translation scheme for $(\mathcal{C}_{bnd} \cup \mathcal{C}_{unb})$-structures $\mathfrak{M}$ whose reducts $\mathfrak{M}|_{\mathcal{C}_{bnd}}$ have tree-width at most k. In order that α and β be mapped to an $\mathrm{MSO}(\mathcal{D}_{bnd})$-sentence and a $C^2(\mathcal{C}_{unb})$-sentence respectively, we need that the translation scheme satisfy some additional properties.

Lemma 14. *Let* $\mathcal{C}_{bnd}$ *and* $\mathcal{C}_{unb}$ *be vocabularies such that* $\mathcal{C}_{bnd} \cap \mathcal{C}_{unb}$ *only contains unary relation symbols. There exist the following effectively computable objects: (1) a vocabulary* $\mathcal{D}_{bnd}$ *consisting of the binary relation symbol* s *and unary relation symbols only, (2) a translation scheme* $\mathbf{tr} = \langle \phi, \psi_C : C \in \mathcal{C}_{bnd} \cup \mathcal{C}_{unb} \rangle$ *for* $\mathcal{C}_{bnd} \cup \mathcal{C}_{unb}$ *over* $\mathcal{D}_{bnd} \cup \mathcal{C}_{unb}$*, and (3) an* $\mathrm{MSO}(\mathcal{D}_{bnd})$*-sentence dom, such that:*

(a) ϕ *is quantifier-free over* $\mathcal{D}_{bnd}$*,*
(b) For every relation symbol $C \in \mathcal{C}_{unb}$*,* ψ_C *is quantifier-free.*
(c) For every relation symbol $C \in \mathcal{C}_{bnd}$*,* ψ_C *is an* $\mathrm{MSO}(\mathcal{D}_{bnd})$*-formula.*
(d) Let $\mathscr{K}$ *be the class of* $(\mathcal{D}_{bnd} \cup \mathcal{C}_{unb})$*-structures in which* s *is interpreted as a binary tree and which satisfy dom. The image of* $\mathscr{K}$ *under* $\mathbf{tr}^\star$ *is exactly the class of* $(\mathcal{C}_{bnd} \cup \mathcal{C}_{unb})$*-structures* $\mathfrak{M}$ *such that* $\mathfrak{M}|_{\mathcal{C}_{bnd}}$ *has tree-width at most* k*.*

The proof of Lemma 14 is technically similar to the discussion in [7]. We include the proof in the Appendix 8.2 for completeness.

We are now ready to prove Lemma 2. By Lemma 14(d) and Lemma 12, statement (i) in Lemma 2 holds iff there is a $(\mathcal{D}_{bnd} \cup \mathcal{C}_{unb})$-structure $\mathfrak{N}$ such that $s^{\mathfrak{N}}$ is a binary tree and $\mathfrak{N} \models dom \wedge \mathbf{tr}^\sharp(\alpha) \wedge \mathbf{tr}^\sharp(\beta)$. Let $\alpha' = dom \wedge \mathbf{tr}^\sharp(\alpha)$ and $\beta' = \mathbf{tr}^\sharp(\beta)$. By Lemma 14(c) and the definition of $\mathbf{tr}^\sharp$, $\alpha' \in \mathrm{MSO}(\mathcal{D}_{bnd})$. Let $\mathbf{tr}|_{\mathcal{C}_{unb}}$ be the translation scheme for $\mathcal{C}_{unb}$ over $\mathcal{D}_{bnd} \cup \mathcal{C}_{unb}$ which agrees with $\mathbf{tr}$ on all formulas, i.e. $\mathbf{tr}|_{\mathcal{C}_{unb}} = \langle \phi, \psi_C : C \in \mathcal{C}_{unb} \rangle$. The image of $\mathbf{tr}|^\star_{\mathcal{C}_{unb}}$ on a structure $\mathfrak{M}$ is the $\mathcal{C}_{unb}$-reduct of the image of $\mathbf{tr}^\star$ on $\mathfrak{M}$. Since $\beta \in C^2(\mathcal{C}_{unb})$, ${\mathbf{tr}|_{\mathcal{C}_{unb}}}^\sharp(\beta)$ is well-defined and ${\mathbf{tr}|_{\mathcal{C}_{unb}}}^\sharp(\beta) = \beta'$. By Lemma 14(a,b), $\mathbf{tr}|_{\mathcal{C}_{unb}}$ is a quantifier-free translation scheme, implying that $\beta' \in C^2(\mathcal{D}_{bnd} \cup \mathcal{C}_{unb})$ by Lemma 13.

---

[5] All translation schemes in this paper are scalar (i.e. non-vectorized). In the notation of [6], a translation scheme is a parameterless non-copying MSO-definition scheme with precondition formula $(x \approx x)$.

6. From MSO to C^2 on Binary Trees

The purpose of this section is to show that, on structures consisting only of a binary tree and additional unary relations, every MSO-sentence can be rewritten to a C^2-sentence which is equi-satisfiable and whose length is linear in the length of the input MSO-sentence. We start by introducing some tools for the literature.

Theorem 4 (Hintikka sentences)**.** *Let $\mathcal{C}$ be a vocabulary. For every $q \in \mathbb{N}$ there is a finite set $\mathscr{HIN}_{\mathcal{C},q}$ of* $\mathrm{MSO}(\mathcal{C})$ *of quantifier rank q such that:*

1. *every $\epsilon \in \mathscr{HIN}_{\mathcal{C},q}$ has a model;*
2. *the conjunction of any two distinct sentences $\epsilon_1, \epsilon_2 \in \mathscr{HIN}_{\mathcal{C},q}$ is not satisfiable;*
3. *every* $\mathrm{MSO}(\mathcal{C})$*-sentence α of quantifier rank at most q is equivalent to exactly one finite disjunction of sentences $\mathscr{HIN}_{\mathcal{C},q}$;*
4. *every $\mathcal{C}$-structure $\mathfrak{A}$ satisfies exactly one sentence* $\mathrm{hin}_{\mathcal{C},q}(\mathfrak{A})$ *of $\mathscr{HIN}_{\mathcal{C},q}$. We may omit $\mathcal{C}$ or q from the subscript when they are clear from the context.*

For a class of $\mathcal{C}$-structures $\mathscr{K}$ an n-ary operation Op over $\mathcal{C}$-structures is called *smooth over $\mathscr{K}$*, if for all $\mathfrak{A}_1, \ldots, \mathfrak{A}_n \in \mathscr{K}$, $\mathrm{hin}_{\mathcal{C},q}(Op(\mathfrak{A}_1, \ldots, \mathfrak{A}_n))$ depends only on $\mathrm{hin}_{\mathcal{C},q}(\mathfrak{A}_i)$: $i \in [n]$ and this dependence is computable[6]. We omit *"over $\mathscr{K}$"* when $\mathscr{K}$ consists of all $\mathcal{C}$-structures.

Theorem 5 (Smoothness)**.** *(1) The disjoint union is smooth.*

(2) For every quantifier-free translation scheme $\mathbf{t}$, the operation $\mathbf{t}^\star$ is smooth.

(3) Let $\mathfrak{T}_1 \overset{\rightarrow}{\circ} \mathfrak{T}_2$ denote the operation which augments the disjoint union of $\mathfrak{T}_1$ and $\mathfrak{T}_2$ by adding an edge from the root of tree $\mathfrak{T}_1$ to the root of tree $\mathfrak{T}_2$. This operation is smooth over labeled trees.[7]

For an in-depth introduction to Hintikka sentences and smoothness and references to proofs see [14, Chapter 3, Theorem 3.3.2] and [19] respectively.

We are now ready for the main lemma of this section.

Let $\mathrm{rt}(x)$ be the sentence $\forall y \neg s(y,x)$. This sentence defines the root of the binary tree s.

Lemma 15. *Let $q \in \mathbb{N}$ and let $\mathcal{C}$ be a vocabulary which consists only of the binary relation symbol s and (possibly) additional unary relation symbols. Let $C_\epsilon : \epsilon \in \mathscr{HIN}_{\mathcal{C},q}$ be new unary relation symbols. Let* $\mathrm{hin}(\mathcal{C},q)$ *be the vocabulary which extends $\mathcal{C}$ with $\{C_\epsilon : \epsilon \in \mathscr{HIN}_{\mathcal{C},q}\}$. There is a computable* $C^2(\mathrm{hin}(\mathcal{C},q))$*-sentence $\Theta^{\mathrm{hin}}_{\mathcal{C},q}$ such that if $\mathfrak{T}_0$ is a $\mathcal{C}$-structure such that $s^{\mathfrak{T}_0}$ is a binary tree:*

---

[6] Smooth operations here are called *effectively smooth* in [19].

[7] We use the smoothness of the disjoint union and quantifier-free transductions. The transduction adds the edge between the roots. Technically, in order for the transduction to be quantifier-free, we need that our trees have a special unary relation symbol root with the natural interpretation as the root of the tree.

(i) *There is an expansion* $\mathfrak{T}_1$ *of* $\mathfrak{T}_0$ *to* $\mathrm{hin}(\mathcal{C}, q)$ *with* $\mathfrak{T}_1 \models \Theta^{\mathrm{hin}}_{\mathcal{C},q}$.

(ii) *For every expansion* $\mathfrak{T}_1$ *of* $\mathfrak{T}_0$ *to* $\mathrm{hin}(\mathcal{C}, q)$ *and for every* $\omega \in \mathrm{MSO}(\mathcal{C})$ *such that* $\mathfrak{T}_1 \models \Theta^{\mathrm{hin}}_{\mathcal{C},q}$ *and* $\mathrm{qr}(\omega) = q$, *the following holds: there exists a* C^2*-sentence* ω_{hin} *such that* $\mathfrak{T}_0 \models \omega$ *iff* $\mathfrak{T}_1 \models \omega_{\mathrm{hin}}$. *The* C^2*-sentence* ω_{hin} *is*

$$\forall x \left(\mathrm{rt}(x) \rightarrow \bigvee_{\epsilon \in \mathscr{H}\mathscr{I}\mathscr{N}_{\mathcal{C},q} :\, \epsilon \models \omega} C_\epsilon(x) \right)$$

The sentence $\Theta^{\mathrm{hin}}_{\mathcal{C},q}$ is defined so that for every $\mathfrak{T}_0$ there is a unique expansion $\mathfrak{T}_1$ such that $\mathfrak{T}_1 \models \Theta^{\mathrm{hin}}_{\mathcal{C},q}$. For every u in the universe T_1 of $\mathfrak{T}_1$, we will have $u \in C_\epsilon^{\mathfrak{T}_1}$ iff the subtree $\mathfrak{T}_u$ of $\mathfrak{T}_0$ whose root is u satisfies $\mathfrak{T}_u \models \epsilon$. Using the smoothness of $\vec{\circ}$, whether an element of $\mathfrak{T}_1$ belongs to $C_\epsilon^{\mathfrak{T}_1}$ depends only on its children. This can be axiomatized in C^2. Lemma 3 follows from Lemma 15 with $q = \mathrm{qr}(\alpha)$, $\mathcal{D} = \mathrm{hin}(\mathcal{C}, q)$, and $\omega_{\mathrm{hin}} = \gamma$.

Appendix 8.3 spells out the proof of Lemma 15.

7. MSO WITH CARDINALITY CONSTRAINTS

$\mathrm{MSO}^{\mathrm{card}}$ denotes the extension of MSO with atomic formulas called *cardinality constraints* $\sum_{i=1}^{r} |X_i| < \sum_{i=1}^{t} |Y_i|$, where the X_i and Y_i are MSO variables, and $|X|$ denotes the cardinality of X. Let WS1S ($\mathrm{WS1S}^{\mathrm{card}}$) be the weak monadic second order theory (with cardinality constraints) of the structure $\langle \mathbb{N}, +1, < \rangle$. Let $\mathrm{MSO}^{\exists\mathrm{card}} \subseteq \mathrm{MSO}^{\mathrm{card}}$ be the set of sentences ρ such that (1) ρ is of the form $\rho = \exists X_1 \cdots \exists X_m \omega$, and (2) only the $X_1, \ldots, X_m$ participate in cardinality constraints.

Theorem 6. *Given a sentence* $\rho \in \mathrm{MSO}^{\exists\mathrm{card}}$, *it is decidable (A) whether* $\langle \mathbb{N}, +1, < \rangle \models \rho$, *and (B) whether* ρ *is satisfiable by a finite structure of bounded tree-width.*

This theorem follows from Theorem 1. The main observation needed for (B) is that cardinality constraints can be expressed in terms of injective functions, which are axiomatizable in C^2. (A) is reducible to (B). The main observations for (A) are:

1. that $X_1, \ldots, X_m$ are contained in the substructure $\mathfrak{A}_1$ of $\langle \mathbb{N}, +1, < \rangle$ generated by $\{0, \ldots, \ell\}$ for some $\ell \in \mathbb{N}$,
2. that substructure $\mathfrak{A}_2$ of $\langle \mathbb{N}, +1, < \rangle$ generated by $\mathbb{N} - \{0, \ldots, \ell\}$ is isomorphic to $\langle \mathbb{N}, +1, < \rangle$, and therefore $\mathfrak{A}_2$ and $\langle \mathbb{N}, +1, < \rangle$ have the same weak monadic second order theory,
3. that the weak monadic second order theory of $\langle \mathbb{N}, +1, < \rangle$ is decidable,
4. and that $\langle \mathbb{N}, +1, < \rangle$ is a transduction $\mathbf{t}$ of $\mathfrak{A}_1 \sqcup \mathfrak{A}_2$.

It remains to use that $\sqcup$ is smooth, cf. Appendix 8.1.

References

[1] S. Arnborg, J. Lagergren, and D. Seese. Easy problems for tree-decomposable graphs. *Journal of Algorithms*, 12(2):308–340, 1991.

[2] F. Baader, D. Calvanese, D. L. McGuinness, D. Nardi, and P. F. Patel-Schneider, editors. *The Description Logic handbook*. Cambridge University Press, 2003.

[3] D. Calvanese, T. Kotek, M. Šimkus, H. Veith, and F. Zuleger. Shape and content. In *Integrated Formal Methods*, pages 3–17. Springer, 2014.

[4] W. Charatonik and P. Witkowski. Two-variable logic with counting and trees. In *LICS*, pages 73–82. IEEE, 2013.

[5] B. Courcelle. The monadic second-order logic of graphs. i. recognizable sets of finite graphs. *Information and computation*, 85(1):12–75, 1990.

[6] B. Courcelle and J. Engelfriet. *Graph structure and monadic second-order logic: a language-theoretic approach*, volume 138. Cambridge University Press, 2012.

[7] B. Courcelle and J. A. Makowsky. Fusion in relational structures and the verification of monadic second-order properties. *Mathematical Structures in Computer Science*, 12(02):203–235, 2002.

[8] J. Flum and M. Grohe. *Parameterized Complexity Theory*. Texts in Theoretical Computer Science. Springer, 2006.

[9] E. Grädel. On the restraining power of guards. *The Journal of Symbolic Logic*, 64(04):1719–1742, 1999.

[10] E. Grädel and M. Otto. On logics with two variables. *Theoretical computer science*, 224(1):73–113, 1999.

[11] E. Grädel, M. Otto, and E. Rosen. Two-variable logic with counting is decidable. In *LICS*, pages 306–317. IEEE, 1997.

[12] E. Grädel, M. Otto, and E. Rosen. Undecidability results on two-variable logics. In *STACS*, pages 249–260, 1997.

[13] J.G. Henriksen, J. Jensen, M. Jørgensen, N. Klarlund, R. Paige, T. Rauhe, and A. Sandholm. *Mona: Monadic second-order logic in practice*. Springer, 1995.

[14] W. Hodges. Model theory. In *Encyclopedia of Mathematics and its Applications*, volume 42. Cambridge University Press, 1993.

[15] F. Klaedtke and H. Rueß. Monadic second-order logics with cardinalities. In *Automata, Languages and Programming*, pages 681–696. Springer, 2003.

[16] T. Kotek and J. A. Makowsky. Connection matrices and the definability of graph parameters. *Logical Methods in Computer Science*, 10(4), 2014.

[17] L. Lovász. *Large networks and graph limits*, volume 60. AMS, 2012.

[18] P. Madhusudan, P.S. Thiagarajan, and S. Yang. The MSO theory of connectedly communicating processes. In *FSTTCS*, pages 201–212. 2005.

[19] J.A. Makowsky. Algorithmic uses of the Feferman–Vaught theorem. *Annals of Pure and Applied Logic*, 126(1):159–213, 2004.

[20] G. Nelson and D.C. Oppen. Simplification by cooperating decision procedures. *TOPLAS*, 1(2):245–257, 1979.

[21] A. Pnueli. The temporal logic of programs. In *FOCS*, pages 46–57. IEEE, 1977.

[22] I. Pratt-Hartmann. Complexity of the two-variable fragment with counting quantifiers. *Journal of Logic, Language and Information*, 14(3):369–395, 2005.

[23] A.J.A. Robinson and A. Voronkov. *Handbook of automated reasoning*. 2001.

[24] D. Seese. The structure of the models of decidable monadic theories of graphs. *Annals of pure and applied logic*, 53(2):169–195, 1991.

[25] W. Thomas. *Languages, automata, and logic*. Springer, 1997.

[26] M.Y. Vardi. Why is modal logic so robustly decidable? *Descriptive complexity and finite models*, 31:149–184, 1996.

8. Appendix

8.1. **Proof of Theorem 6.** We start with (B).

Let ρ be a $\mathrm{MSO}^{\exists \mathrm{card}}$ sentence, i.e. the outermost block of quantifiers in ρ is existential and only variables from the outermost blocks may appear in cardinality constraints. For simplicity we consider $\rho = \exists X_1 \exists X_2 \omega$ with only two quantifiers in the outermost block. W.l.o.g. the only cardinality constraint in ω is $|X_1| < |X_2|$. By a slight abuse of notation, we sometimes treat X_1 and X_2 as unary relation symbols. Let $\mathcal{C}_{bnd}$ extend the vocabulary of ρ with new unary relation symbols X_1,X_2,W_{img},W_{dom}. Let $\mathcal{C}_{unb}$ extend $\mathcal{C}_{bnd}$ with a new binary relation symbol B. Finite satisfiability of ω by a structure $\mathfrak{M}$ such that $\mathrm{tw}(\mathfrak{M}) \leq k$ can be reduced to finite satisfiability of a sentence $\alpha \wedge \beta$, $\alpha \in \mathrm{MSO}(\mathcal{C}_{bnd})$ and $\beta \in C^2(\mathcal{C}_{unb})$, by a structure $\mathfrak{A}_1$ such that $\mathrm{tw}(\mathfrak{A}_1|_{\mathcal{C}_{bnd}}) \leq k$. Let β be the C^2-sentence

$$\beta = (\mathrm{inj}_{12} \vee \mathrm{inj}_{21}) \wedge \mathrm{dom} \wedge \mathrm{img}$$

where

- inj_{12} expresses that B is an injective function from X_1 to X_2,
- inj_{21} espresses that B is an injective function from X_2 to X_1,
- dom expresses that the domain of B is W_{dom},
- and img expresses that the image of B is W_{img}.

For every $\mathcal{C}_{unb}$-structure $\mathfrak{A}_1$, $|X_1^{\mathfrak{A}_1}| < |X_2^{\mathfrak{A}_1}|$ iff $W_{\mathrm{dom}}^{\mathfrak{A}_1} = X_1^{\mathfrak{A}_1}$ and $X_2^{\mathfrak{A}_1} \backslash W_{\mathrm{img}}^{\mathfrak{A}_1} \neq \emptyset$. Let α be obtained from ω by substituting every $|X_1| < |X_2|$ by

$$\forall x\, (X_1(x) \leftrightarrow W_{\mathrm{dom}}(x)) \wedge \exists x\, (\neg W_{\mathrm{img}}(x) \wedge X_2(x))$$

For any $\mathcal{C}_{bnd}$-structure $\mathfrak{A}_0$ with $\mathrm{tw}(\mathfrak{A}_0) \leq k$, $\mathfrak{A}_0 \models \omega$ iff there is an expansion $\mathfrak{A}_1$ of $\mathfrak{A}_0$ such that $\mathfrak{A}_1 \models \alpha \wedge \beta$. The treatment of other cardinality constraints $\sum_{i=1}^{r} |X_i| < \sum_{i=1}^{t} |Y_i|$ is similar; it is helpful to assume w.l.o.g. that the X_i and the Y_i are disjoint.

Now we turn to (A). The unary function $+1$ is the *successor relation* of $\mathbb{N}$ and interprets the binary relation symbol suc. The binary relation $<$ is the *natural order relation* on $\mathbb{N}$. In the proof of (A) we will use the theory of Hintikka sentences as presented in Section 6 with one caveat, namely that instead of restricting to finite structures, we allow arbitrary structures. Theorems 4 and 5 hold for arbitrary structures. Theorems 4 guarantees the existence of a set $\mathscr{HIN}_{\mathcal{C},q}^{\mathrm{arb}}$ analogously to $\mathscr{HIN}_{\mathcal{C},q}$ for arbitrary structures. For every $\mathcal{C}$-structure $\mathfrak{A}$, Theorem 4 guarantees the existence of a sentence $\mathrm{hin}_{\mathcal{C},q}^{\mathrm{arb}}(\mathfrak{A})$ of $\mathscr{HIN}_{\mathcal{C},q}^{\mathrm{arb}}$ analogously to $\mathrm{hin}_{\mathcal{C},q}(\mathfrak{A})$. For the rest of the section, we omit the superscript arb to simplify notation.

Consider $\rho = \exists X_1 \exists X_2 \omega$ for the vocabulary $\mathcal{C}_{\mathbb{N}}$ of $\langle \mathbb{N}, +1, < \rangle$. Let $\alpha \in \mathrm{MSO}(\mathcal{C}_{bnd})$ and $\beta \in C^2(\mathcal{C}_{unb})$ be as discussed in the proof of (B) above. The following are equivalent:

1. $\langle \mathbb{N}, +1, < \rangle \models \rho$
2. There are *finite* unary relations U_1 and U_2 such that $\langle \mathbb{N}, +1, <, U_1, U_2 \rangle \models \omega$. U_1 and U_2 interpret X_1 and X_2 respectively.
3. There are *finite* unary relations U_1 and U_2 and an expansion $\mathfrak{A}$ of $\langle \mathbb{N}, +1, <, U_1, U_2 \rangle$ such that $\mathfrak{A} \models \alpha \wedge \beta$. $\mathfrak{A}$ expands $\langle \mathbb{N}, +1, <, U_1, U_2 \rangle$ with interpretations of the symbols in $\mathcal{D}_{unb} = \{B, W_{\mathrm{dom}}, W_{\mathrm{img}}\}$.
4. $\rho' = \exists X_1 \exists X_2 (\alpha \wedge \beta)$ is satisfiable by an expansion of $\langle \mathbb{N}, +1, < \rangle$ with interpretations for the symbols of $\mathcal{D}_{unb}$.

We have *1. iff 2.* and *3. iff 4.* by the semantics of $\exists X_1 \exists X_2$ in weak monadic second order logic. We have *2. iff 3.* similarly to the discussion of α and β in the proof of (A) above. The rest of the proof is devoted to proving that 4. is decidable.

Observe that by the definition of β, and in weak monadic second order X_1 and X_2 are quantified to be finite sets, B is axiomatized to be a function with finite domain, and W_{dom} and W_{img} are finite. Hence models $\mathfrak{A}$ of ρ' can be decomposed into a finite part containing $B^{\mathfrak{A}}$, and an infinite part isomorphic to an expansion of $\langle \mathbb{N}, +1, < \rangle$ in which the symbols of $\mathcal{D}_{unb}$ as are interpreted as empty sets. We will use a similar decomposition, but first we want to move from the structure $\langle \mathbb{N}, +1, < \rangle$ and its expansions to $\langle \mathbb{N}, +1 \rangle$ and its expansions. There is a translation scheme $\mathbf{t}_<$ such that for every structure $\mathfrak{A} = \langle \mathbb{N}, +1, B^{\mathfrak{A}}, W^{\mathfrak{A}}_{\mathrm{dom}}, W^{\mathfrak{A}}_{\mathrm{img}} \rangle$, $\mathbf{t}^{\star}_<(\langle \mathfrak{A} \rangle) = \langle \mathfrak{A}, < \rangle$, where $\langle \mathfrak{A}, < \rangle$ is the expansion of $\mathfrak{A}$ with $<$. This is true since $<$ is MSO definable from $+1$. We have that ρ' is satisfied by an expansion of $\langle \mathbb{N}, +1, < \rangle$ iff $\mathbf{t}^{\sharp}_<(\rho')$ is satisfied by the same expansion of $\langle \mathbb{N}, +1 \rangle$.

Now we turn to the decomposition of models of $\mathbf{t}^{\sharp}_<(\rho')$ into a finite part containing $B^{\mathfrak{A}}$, and an infinite part isomorphic to a $\mathcal{D}_{unb}$-expansion of $\langle \mathbb{N}, +1 \rangle$. Let $\mathcal{D}_2 \supseteq \mathcal{D}_{unb}$ be the vocabulary of $\mathbf{t}^{\sharp}_<(\rho')$. For every $\mathcal{D}_2$-expansion $\mathfrak{P}$ of $\langle \mathbb{N}, +1 \rangle$ and every $n \in \mathbb{N}$, let $\mathfrak{P}_{1,n}$ and $\mathfrak{P}_{n,\infty}$ be the substructures of $\mathfrak{P}$ generated by $[n]$ and $\mathbb{N} \backslash [n-1]$ respectively. There is a translation scheme $\mathbf{u}$ such that $\mathbf{u}^{\star}(\mathfrak{P}_{1,n} \sqcup \mathfrak{P}_{n+1,\infty}) = \mathfrak{P}$ if $B^{\mathfrak{P}} \subseteq [n] \times [n]$. $\mathbf{u}$ existentially quantifies the set $[n]$ (which is the only non-empty finite set closed under suc and its inverse), 1 and n (as the first and last elements of $[n]$) and $n+1$ (as the only element without a suc-predecessor except for 1) and adds the edge $(n, n+1)$ to suc. We have $\mathfrak{P} \models \mathbf{t}^{\sharp}_<(\rho')$ iff $\mathfrak{P}_{1,n} \sqcup \mathfrak{P}_{n+1,\infty} \models \mathbf{u}^{\sharp}(\mathbf{t}^{\sharp}_<(\rho'))$. Note that the vocabulary of $\mathbf{u}^{\sharp}(\mathbf{t}^{\sharp}_<(\rho'))$ is $\mathcal{D}_2$. Let q be the quantifier rank of $\mathbf{u}^{\sharp}(\mathbf{t}^{\sharp}_<(\rho'))$.

The Hintikka sentence $\mathrm{hin}(\mathfrak{P}_{n+1,\infty})$ of quantifier rank q of $\mathfrak{P}_{n+1,\infty}$ is uniquely defined since $\mathfrak{P}_{n+1,\infty}$ is isomorphic to the expansion of $\langle \mathbb{N}, +1 \rangle$ with empty sets. Moreover, $\mathrm{hin}(\mathfrak{P}_{n+1,\infty})$ is computable using that the theory of $\langle \mathbb{N}, +1 \rangle$ is decidable. Hence, by the smoothness of the disjoint union, for every Hintikka sentence $\epsilon \in \mathscr{HIN}_{\mathcal{D}_2,q}$ there is a computable set $E_\epsilon \subseteq \mathscr{HIN}_{\mathcal{D}_2,q}$ such that $\mathrm{hin}(\mathfrak{P}_{1,n} \sqcup \mathfrak{P}_{n+1,\infty}) = \epsilon$ iff $\mathrm{hin}(P_{1,n}) \in E_\epsilon$. Then $\mathfrak{P}_{1,n} \sqcup \mathfrak{P}_{n+1,\infty} \models \mathbf{u}^{\sharp}(\mathbf{t}^{\sharp}_<(\rho'))$ iff $\mathfrak{P}_{1,n}$ satisfies the sentence $\bigvee_{(\epsilon,\epsilon')} \epsilon'$, where the $\bigvee_{(\epsilon,\epsilon')}$ ranges over pairs (ϵ, ϵ') such that

1. $\epsilon \in \mathscr{HIN}_{\mathcal{D}_2,q}$,
2. $\epsilon \models \mathbf{u}^{\sharp}(\mathbf{t}^{\sharp}_<(\rho'))$, and
3. $\epsilon' \in E_\epsilon$.

Hence, ρ' is satisfiable by an expansion of $\langle \mathbb{N}, +1, < \rangle$ iff $\bigvee_{(\epsilon,\epsilon')} \epsilon'$ is satisfiable by a finite structure in which suc is interpreted as a successor relation (i.e., as a simple directed path on the whole universe). Let suc-rel be the weak MSO-sentence such that the interpretation of suc is a successor relation. By Theorem 6(B), it is decidable whether, $\bigvee_{(\epsilon,\epsilon')} \epsilon' \wedge$ suc-rel is finitely satisfiable using that the class of simple directed paths annotated with unary relations has tree-width 1.

Remark 1. *While we assumed for simplicity in (B) that X_1 and X_2 range over subsets of the universe, it is not hard to extend the proof to the case that X_1 and X_2 are guarded second order variables which range over subsets of any relation in the structure. This is true since we can use the translation scheme* $\mathbf{tr}$ *from Lemma 14 to obtain the structures of tree-width at most k as the image of* $\mathbf{tr}^{\star}$ *of labeled trees; X_1 and X_2 then translate naturally to monadic second order variables.*

8.2. **Proof of Lemma 14.** This appendix is devoted to the proof of Lemma 14. Appendix 8.2 introduces a tree encoding of structures of tree-width at most k based on *aligned k-trees*. Appendix 8.2 introduces a translation scheme such that the image of the induced transduction on an MSO-definable class of labeled binary trees is the class of structures of tree-width at most k. This is done based on translation schemes introduced in Appendix 8.2. Finally, the proof of Lemma 14 is given in Subsection 8.2.

Note that the notion of *aligned k-trees* is quite similar to the notion of *oriented k-trees* defined in Section 2. In fact, we could have replaced aligned k-trees with oriented k-trees here, or replaced oriented k-trees with aligned k-trees throughout the paper. However, it simplifies the proofs to have two notions.

Aligned k-tree encodings and (decorated) aligned k-trees. An aligned k-tree encoding $\mathfrak{T}$ is a binary tree whose vertices are labeled by certain unary relations. All structures of bounded tree-width can be obtained by applying transductions to aligned k-tree encodings with additional vertex annotations in three steps:

Step A	Aligned k-tree encoding	$\xrightarrow{\textbf{interpret}}$	Decorated aligned k-tree
Step B	Decorated aligned k-tree	$\xrightarrow{\textbf{undecorate}}$	Aligned k-tree
Step C	Aligned k-tree	$\xrightarrow{\textbf{structurize}_{\mathcal{C}}}$	$\mathcal{C}$-structure of tw $\leq k$

interpret, **undecorate** and **structurize**$_{\mathcal{C}}$ (shorthand **i**, **u** and $\mathbf{s}_{\mathcal{C}}$) are unary operations on structures given in terms of translation schemes (see Section 8.2).

Two key properties of our encoding:

(a) Using that the number of edges in a k-tree is at most $k \cdot$ (number of vertices), our (decorated) aligned k-trees have functions $R_1, \ldots, R_k$ rather than an edge relation.

(b) The elements of the universe of a structure appear as elements in the aligned k-tree, in the decorated aligned k-tree, and in the aligned k-tree encoding. Aligned k-tree encodings and decorated aligned k-trees have additional auxiliary vertices which are eliminated by **undecorate**.

Let $\mathcal{V}$ be the vocabulary containing the binary relation symbol s, and the unary relation symbols $\mathrm{root}, \mathrm{Label}_1, \ldots, \mathrm{Label}_k, \mathrm{Label}_{\mathrm{blank}}$.

An **aligned k-tree encoding** is a $\mathcal{V}$-structure $\mathfrak{T}$ with universe T such that (i) $(T, s^{\mathfrak{T}})$ is a directed tree (i.e., an acyclic directed graph in which all vertices have in-degree 1 except exactly one.), (ii) every vertex in $(T, s^{\mathfrak{T}})$ has out-degree at most 2, (iii) $\mathrm{Label}_1^{\mathfrak{T}}, \ldots, \mathrm{Label}_k^{\mathfrak{T}}, \mathrm{Label}_{\mathrm{blank}}^{\mathfrak{T}}$ form a partition of T, (iv) $\mathrm{root}^{\mathfrak{T}}$ denotes the unary relation containing only the unique vertex with in-degree 0 in $(T, s^{\mathfrak{T}})$, and (v) all the children of vertices in $T \backslash \mathrm{Label}_{\mathrm{blank}}^{\mathfrak{T}}$ belong to $\mathrm{Label}_{\mathrm{blank}}^{\mathfrak{T}}$. Fig. 3(a) shows an aligned 3-tree encoding. The edges represent s. The small black circles represent $\mathrm{Label}_{\mathrm{blank}}$, the larger red circles represent Label_1, the squares represent Label_2, and the diamonds represent Label_3.

Lemma 16. *There is a C^2-sentence* enc *such that for every $\mathcal{V}$-structure $\mathfrak{M}$ in which $s^{\mathfrak{M}}$ is a binary tree, $\mathfrak{M} \models$* enc *iff $\mathfrak{M}$ is an aligned k-tree encoding.*

Proof. Being an aligned k-tree encoding is expressible in C^2 for structures in which s is interpreted as a binary tree. Below ξ_X refers to requirement X in the definition of an aligned k-tree encoding. ξ_i and ξ_{ii} are already taken care of since s is a binary



FIGURE 3. **(a)** An aligned 3-tree encoding. **(b)** The decorated aligned 3-tree obtained from (a) by applying $\mathbf{i}^\star$. R_1, R_2 and R_3 are denoted by the dashed arrows, the solid arrows and the (single) thick arrow, respectively. **(c)** The aligned 3-tree obtained from (b) by applying $\mathbf{u}^\star$. **(d)** A aligned 3-tree $\mathfrak{A}$ expanding (c) with unary relations annotating its elements. $\mathfrak{A}$ encodes a directed graph $G = (V, E)$. The unary relations are $U_j^{\mathfrak{A}}$ and $U_{\mathrm{inv},j}^{\mathfrak{A}}$ for each $j \in \{1, 2, 3\}$. They encode, for each element m of $\mathfrak{A}$, whether the edge $(m, R_j(m))$ respectively $(R_j(m), m)$ belongs to E. **(e)** The directed graph obtained from $\mathfrak{A}$ in (d) by applying $\mathbf{s}^\star$. $\mathrm{Gaif}(G)$ has tree-width 2.

tree. enc is given as follows:

$$\begin{aligned}
\mathrm{enc} &= \xi_{iii} \wedge \xi_{iv} \wedge \xi_v \\
\xi_{iii} &= \forall x \left(\mathrm{Label}_{\mathrm{blank}}(x) \vee \bigvee_{i=1}^{k} \mathrm{Label}_i(x) \right) \wedge \\
&\quad\ \forall x \left(\bigvee_{i=1}^{k} \mathrm{Label}_i(x) \leftrightarrow \neg \mathrm{Label}_{\mathrm{blank}}(x) \right) \wedge \\
&\quad\ \forall x \left(\bigwedge_{i \neq j} (\neg \mathrm{Label}_i(x) \vee \neg \mathrm{Label}_j(x)) \right) \\
\xi_{\mathrm{iv}} &= \forall x \left(\mathrm{root}(x) \leftrightarrow \forall y \neg s(y, x) \right) \\
\xi_v &= \forall x \forall y \left(s(x, y) \wedge \neg \mathrm{Label}_{\mathrm{blank}}(x) \rightarrow \mathrm{Label}_{\mathrm{blank}}(y) \right)
\end{aligned}$$

□

A **decorated aligned k-tree** $\mathfrak{A}$ is an expansion of an aligned k-tree encoding $\mathfrak{T}$ to $\mathcal{R} \cup \mathcal{R}$. For every $j \in [k]$, $R_j^{\mathfrak{A}}$ contains all pairs $(v, u) \in (T \backslash (\mathrm{Label}_j^{\mathfrak{T}} \cup \mathrm{Label}_{\mathrm{blank}}^{\mathfrak{T}})) \times \mathrm{Label}_j^{\mathfrak{T}}$ such that there is a directed path P from u to v in $(T, s^{\mathfrak{T}})$ which does not intersect with $\mathrm{Label}_j^{\mathfrak{T}}$ except on u, i.e. $\mathrm{Label}_j^{\mathfrak{T}} \cap (P \backslash \{u\}) = \emptyset$; moreover, for every u, if $R_j^{\mathfrak{A}}$ does not contain any pair (u, v), then $R_j^{\mathfrak{A}}$ contains additionally the self loop (u, u). Observe that the relation $R_j^{\mathfrak{A}}$ is a total function. Fig. 3(b) shows the decorated aligned 3-tree obtained from (a). The new edges represent R_1, R_2, and R_3: dashed edges represent R_1, solid edges represent R_2, and the single thick edge (from a square to a diamond) represents R_3.

A **aligned** k**-tree**[8] is the $\mathcal{R}$-reduct of the substructure of a decorated aligned k-tree $\mathfrak{A}_0$ generated by $\mathrm{Label}_1^{\mathfrak{A}_0} \cup \cdots \cup \mathrm{Label}_k^{\mathfrak{A}_0}$. Fig. 3(c) shows the aligned 3-tree obtained from (b). The vertices of (c) are not labeled by any unary relation.

Lemma 17.

(1) For every aligned k-tree $\mathfrak{A}$, $\mathrm{Gaif}(\mathfrak{A})$ *is a partial $(k-1)$-tree.*

(2) For every partial $(k-1)$-tree G there is an aligned k-tree $\mathfrak{A}$ such that G is a subgraph of $\mathrm{Gaif}(\mathfrak{A})$.

Proof. For the proof of Lemma 17(1) we define an aligned k-tree encoding $\mathfrak{T}$ to be **perfect** if (1) there are $w_1, p_1, w_2, \ldots, p_{k-1}, w_k \in T$ such that $w_1, p_1, w_2, \ldots, p_{k-1}, w_k$ is a directed path, (2) $\mathrm{root}^{\mathfrak{T}} = \{w_1\}$, (3) $w_i \in \mathrm{Label}_i^{\mathfrak{T}_0}$ of all $i \in [k]$, (4) $p_i \in \mathrm{Label}_{\mathrm{blank}}$ for all $i \in [k-1]$, and (5) the out-degree of each of vertex in the path is exactly 1. If $\mathfrak{T}$ is a perfect aligned k-tree encoding, then $\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}))$ is a perfect aligned k-tree. We have:

($\diamond$) For every perfect aligned k-tree $\mathfrak{A}$, $\mathrm{Gaif}(\mathfrak{A})$ is a $(k-1)$-tree.

($\diamond$) is proved by induction on the number of vertices in a perfect aligned k-tree encoding $\mathfrak{T}$ such that $\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})) = \mathfrak{A}$. Lemma 17(1) follows from ($\diamond$) by building a perfect aligned k-tree such that $\mathfrak{A}$ is its substructure.

First we prove:

($\diamond$) For every perfect aligned k-tree $\mathfrak{A}$, $\mathrm{Gaif}(\mathfrak{A})$ is a $(k-1)$-tree.

Let $\mathfrak{T}$ be a perfect aligned k-tree encoding such that $\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})) = \mathfrak{A}$. We prove the claim by induction on the number of vertices in $\mathfrak{T}$.

- In the base case, $\mathfrak{T}$ consists only of the vertices $w_1, p_1, \ldots, p_{k-1}, w_k$, and $\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}))$ is a k-clique, which is a $(k-1)$-tree.
- Let $\mathfrak{T}$ be a perfect aligned $(k-1)$-tree encoding. Let $\mathfrak{T}_0$ be obtained from $\mathfrak{T}$ by removing a leaf v. By the induction hypothesis, $\mathrm{Gaif}(\mathbf{i}^\star(\mathfrak{T}_0))$ is a $(k-1)$-tree. If $v \in \mathrm{Label}_{\mathrm{blank}}^{\mathfrak{T}}$, then $\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}_0)) = \mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}))$. Otherwise, let $v \in \mathrm{Label}_j^{\mathfrak{T}}$. We denote by $B = \{u_i : i \in [k]\backslash\{j\}\}$ the set of elements such that $u_i \in \mathrm{Label}_i$ and there is a path P_i from u_i to v in $\mathfrak{T}$ satisfying the following property: P_i does not visit any vertex in Label_i except u_i. Note that for all $i \in [k]\backslash\{j\}$ u_i exists, and hence $|B| = k-1$, using that $\mathfrak{T}_0$ is perfect. By the choice of B and since v is a leaf, B is the set of neighbors of v in $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})))$. It remains to show that B is a $(k-1)$-clique to get that $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})))$ is a $(k-1)$-tree. Let $i_1, i_2 \in [k]\backslash\{j\}$ with $i_1 \neq i_2$. Since $(T, s^{\mathfrak{T}})$ is a directed tree, either $P_{i_1} \subsetneq P_{i_2}$ or $P_{i_2} \subsetneq P_{i_1}$. W.l.o.g. let $P_{i_1} \subsetneq P_{i_2}$, so P_{i_2} is a path from u_{i_2} to v which visits u_{i_1}. Let $P_{21} \subsetneq P_{i_2}$ be the path from u_{i_2} to u_{i_1}. By the choice of B, there is no vertex in P_{21} which belongs to Label_{i_2} except u_{i_2}. Hence, $(u_{i_1}, u_{i_2}) \in R_{i_2}^{\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}))}$, i.e. there is an edge between u_{i_1} and u_{i_2} in $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})))$. We get that B is a $(k-1)$-clique and $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})))$ is a $(k-1)$-tree.

Now we are ready to prove Lemma 17(1) and Lemma 17(2).

(1) Let $\mathfrak{T}$ be an aligned k-tree encoding such that $\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})) = \mathfrak{A}$. Let $\mathfrak{T}_0$ be obtained from the directed path $w_1, p_1, \ldots, p_{k-1}, w_k$ with $w_i \in \mathrm{Label}_i^{\mathfrak{T}_0}$ for

---

[8] A small but important note is that the definition of aligned k-trees used throughout this appendix deviates slightly from the one given in Section 2 and used elsewhere in the paper.

each $i \in [k]$, $p_i \in \mathrm{Label}_{\mathrm{blank}}^{\mathfrak{T}_0}$ for each $i \in [k-1]$, and $\mathrm{root}^{\mathfrak{T}} = \{w_1\}$ by attaching $\mathfrak{T}$ as a child of w_k. By construction, $\mathfrak{T}_0$ is perfect. By ($\diamond$) we have that $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}_0)))$ is a $(k-1)$-tree. Deleting the vertices $w_1, p_1, \ldots, p_{k-1}, w_k$ and the edges incident to them from $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}_0)))$ we get $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})))$, implying that $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T})))$ is a partial k-tree.

(2) We prove this by induction on the construction of $(k-1)$-trees.
   (a) Let G be a $(k-1)$-clique with vertices $w_1, \ldots, w_{k-1}$. Let $\mathfrak{T}_{base}$ be the directed path on the vertices $w_1, p_1, \ldots, p_{k-2}, w_{k-1}$ such that $w_i \in \mathrm{Label}_i^{\mathfrak{T}_{base}}$ for all $i \in [k-1]$ and $p_i \in \mathrm{Label}_{\mathrm{blank}}^{T_{base}}$ for all $i \in [k-2]$. We have that $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}_{base}))) = G$.
   (b) Let G_0 be a $(k-1)$-tree and let $\mathfrak{T}_0$ be an aligned k-tree encoding such that $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}_0))) = G_0$. Let $C = \{u_1, \ldots, u_{k-1}\}$ be a $(k-1)$-clique in G_0. Let G be the graph obtained from G_0 by adding a new vertex v as well as edges between v and each of the vertices in C. Since C is a $(k-1)$-clique in G_0, for every distinct $i_1, i_2 \in [k-1]$ there is a directed path in $\mathfrak{T}_0$ between from u_{i_1} to u_{i_2} or vice versa. As a consequence, there is a path P in $\mathfrak{T}_0$ from the root to some u_ℓ such that $u_1, \ldots, u_{k-1}$ all occur on P. For every $t \in [k-1]$ and $i \in [k]$ such that $u_t \in \mathrm{Label}_i^{\mathfrak{T}_0}$, u_t is the last vertex in P to belong to $\mathrm{Label}_i^{\mathfrak{T}_0}$. Since $|C| = k-1$, there is $j \in [k]$ such that $\mathrm{Label}_j^{\mathfrak{T}_0} \cap C = \emptyset$. Since C is a clique in $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}_0)))$, it must be the case that $|\mathrm{Label}_i^{\mathfrak{T}_0} \cap C| = 1$ for all $i \in [k] \backslash \{j\}$. If u_ℓ has two children, let $\mathfrak{T}$ be a subtree of $\mathfrak{T}_0$ whose root is a child of u_ℓ. Let $\mathfrak{T}$ be obtained from $\mathfrak{T}_0$ by attaching a new child vertex $v_{\mathrm{blank}} \in \mathrm{Label}_{\mathrm{blank}}^{\mathfrak{T}}$ to u_ℓ, moving $\mathfrak{T}$ to be a child of v_{blank}, and adding v as a child of v_{blank}. If u_ℓ is a leaf or has one child in $\mathfrak{T}_0$, let $\mathfrak{T}$ be obtained from $\mathfrak{T}_0$ by attaching v as a child of u_ℓ. Setting v to belong to $\mathrm{Label}_j^{\mathfrak{T}}$, we get $\mathrm{Gaif}(\mathbf{u}^\star(\mathbf{i}^\star(\mathfrak{T}))) = G$.

□

The translation schemes interpret, undecorate and structurize$_{\mathcal{C}}$. There are translation schemes **interpret** (shorthand **i**) for $\mathcal{R} \cup \mathcal{V}$ over $\mathcal{V}$ and **undecorate** (shorthand **u**) for $\mathcal{R}$ over $\mathcal{R} \cup \mathcal{V}$ which take an aligned k-tree encoding to its decorated aligned k-tree respectively a decorated aligned k-tree to its aligned k-tree. The induced transductions $\mathbf{i}^\star$ and $\mathbf{u}^\star$ are surjective with respect to the classes of decorated aligned k-trees resp. aligned k-trees.

For every vocabulary $\mathcal{C}$ there is a translation scheme **structurize**$_{\mathcal{C}}$ (shorthand $\mathbf{s}_{\mathcal{C}}$) which takes an aligned k-tree whose elements are annotated with unary relations to a $\mathcal{C}$-structure. The induced transduction $\mathbf{s}_{\mathcal{C}}$ is surjective with respect to the class of $\mathcal{C}$-structures of tree-width less than k. The binary relations of such structures are encoded inside aligned k-trees by unary relations on the sources of R_j edges, using that the R_j are functions. The vocabulary consisting of these new unary relation symbols as well as the unary relation symbols of $\mathcal{C}$ is denoted by $\mathcal{N}_{\mathcal{C}}$. $\mathbf{s}_{\mathcal{C}}$ is a translation scheme for $\mathcal{C}$ over $\mathcal{R} \cup \mathcal{N}_{\mathcal{C}}$. Fig. 3(d) shows an aligned k-tree annotated with the unary relations of $\mathcal{N}_{\mathcal{C}}$ for $\mathcal{C} = \langle E \rangle$, where E is a binary relation symbol. Fig. 3(e) shows the directed graph obtained from (d) by applying $\mathbf{s}_{\mathcal{C}}$.

Lemma 18.

A.: *For every decorated aligned k-tree $\mathfrak{A}$, $\mathbf{i}^\star(\mathfrak{A}|_{\mathcal{V}}) = \mathfrak{A}$.*

B.: *For every aligned k-tree $\mathfrak{A}$ obtained from a decorated aligned k-tree $\mathfrak{A}_0$, $\mathbf{u}^\star(\mathfrak{A}_0) = \mathfrak{A}$.*

C.: *There is a vocabulary $\mathcal{N}_\mathcal{C}$ consisting of unary relation symbols only, such that for every $\mathcal{C}$-structure $\mathfrak{M}$, $\mathfrak{M}$ has tree-width at most k iff there is an aligned k-tree $\mathfrak{A}_{ori}$ and an expansion $\mathfrak{A}$ of $\mathfrak{A}_{ori}$ to $\mathcal{N}_\mathcal{C} \cup \mathcal{R}$ such that $\mathbf{s}^\star_\mathcal{C}(\mathfrak{A}) = \mathfrak{M}$.*

Lemma 2 follows from Lemma 18, Lemma 16, and Lemma 12 with $\mathcal{D}_{bnd} = \mathcal{N}_{\mathcal{C}_{bnd}} \cup \mathcal{V}$, $\alpha' = \mathrm{enc} \land \mathbf{i}^\sharp(\mathbf{u}^\sharp(\mathbf{s}^\sharp_\mathcal{C}(\alpha)))$, and $\beta' = \mathbf{i}^\sharp(\mathbf{u}^\sharp(\mathbf{s}^\sharp_\mathcal{C}(\beta)))$.

Proof of Lemma 18(A). $\mathbf{i} = \langle \varphi, \psi_C : C \in \mathcal{R} \rangle$ is given as follows:

1. The universe A of $\mathfrak{A}$ is T, i.e. $\varphi(x) = (x \approx x)$.
2. The relations from $\mathcal{V}$ are copied without change, i.e. $\psi_s(x, y) = s(x, y)$, and $\psi_{\mathrm{Label}}(x) = \mathrm{Label}(x)$ for each $\mathrm{Label} \in \{\mathrm{Label}_1, \ldots, \mathrm{Label}_k, \mathrm{Label}_{\mathrm{blank}}\}$.
3. $\psi_{R_j}(x, y) = \varphi(x) \land \neg\mathrm{Label}_j(x) \land \mathrm{Label}_j(y) \land \forall z\, (z \not\approx y \to \neg\mathrm{Label}_j(z)) \land (\mathrm{dpath}(x, y))$ where $\mathrm{dpath}(x, y)$ says that there is a subset P of s which is a directed simple path from x to y:

$$\begin{aligned}
\mathrm{dpath}(x,y) &= \exists P \subseteq s \Big[\forall z\,(in_{\leq 1}(P,x)) \land in_0(P,x) \land \\
&\qquad\qquad \forall z\,(in_{\leq 1}(P,y)) \land out_0(P,y) \land \mathrm{reach}(P,x,y)\Big] \\
\mathrm{reach}(P,x,y) &= \forall X \subseteq V \Bigg((X(x) \land \neg X(y)) \to \\
&\qquad\qquad (\exists z_1 \exists z_2\, (X(z_1) \land \neg X(z_2) \land P(z_1,z_2))) \Bigg) \\
in_0(z_1,P) &= \forall z_2\, (\neg P(z_2,z_1)) \\
out_0(z_1,P) &= \forall z_2\, (\neg P(z_1,z_2)) \\
in_{\leq 1}(z_1,P) &= \exists^{\leq 1} z_2\, (P(z_2,z_1)) \\
out_{\leq 1}(z_1,P) &= \exists^{\leq 1} z_2\, (P(z_1,z_2))
\end{aligned}$$

   $\mathrm{reach}(P, x, y)$ says that y is reachable from x via P edges.

Proof of Lemma 18(B). Let $\mathbf{u} = \langle \varphi, \psi_{R_1}, \ldots, \psi_{R_k} \rangle$ be as follows: $\varphi(x) = \neg\mathrm{Label}_{\mathrm{blank}}(x)$ and for every $i \in [k]$, $\psi_{R_i}(x, y) = R_i(x, y)$.

Proof of Lemma 18(C). We prove the following lemma which spells out the two directions of Lemma 18(C):

Lemma 19. *For every vocabulary $\mathcal{C}$, there is a set $\mathcal{N}_\mathcal{C}$ of unary relation symbols and a translation scheme* $\mathbf{structurize}_\mathcal{C}$ *(shorthand $\mathbf{s}_\mathcal{C}$) for $\mathcal{C}$ over $\mathcal{N}_\mathcal{C} \cup \mathcal{R}$ such that:*

(1) If $\mathfrak{M}$ is a $\mathcal{C}$ structure whose Gaifman graph is a partial $(k-1)$-tree, then there is an expansion $\mathfrak{A}$ of an aligned k-tree to $\mathcal{N}_\mathcal{C} \cup \mathcal{R}$ for which $\mathbf{s}^\star_\mathcal{C}(\mathfrak{A}) = \mathfrak{M}$.

(2) If $\mathfrak{A}$ is a $(\mathcal{N}_\mathcal{C} \cup \mathcal{R})$-structure and $\mathfrak{A}_{red} = \mathfrak{A}|_\mathcal{R}$ is an aligned k-tree, then $\mathrm{Gaif}(\mathbf{s}^\star_\mathcal{C}(\mathfrak{A}))$ *is a partial $(k-1)$-tree.*

For every vocabulary $\mathcal{C}$, let $\mathcal{N}_\mathcal{C}$ be the vocabulary extending the set $\mathrm{un}(\mathcal{C})$ of unary relation symbols in $\mathcal{C}$ by fresh unary relation symbols $U_{B,\mathrm{self}}$, $U_{B,j}$, and $U_{B,\mathrm{inv},j}$ for every binary relation symbol $B \in \mathcal{C}$ and $j \in [k]$. Recall $\mathcal{V}_\mathcal{C} = \mathcal{V} \cup \mathcal{N}_\mathcal{C}$ and $\mathcal{R}_\mathcal{C} = \mathcal{V}_\mathcal{C} \cup \mathcal{R} \cup \mathcal{C}$.

Since the R_i are functions, we use a fixed number of unary relation symbols to encode, for every universe element m_1 such that $(m_1, m_2) \in R_j^\mathfrak{M}$, what other relations of $\mathfrak{M}$ do (m_1, m_2) and (m_2, m_1) belong to. The unary relation symbols are of the form $U_{B,j}$ and $U_{B,\mathrm{inv},j}$, where $j \in [k]$ and B is a binary relation symbol. We use that tuples (m_1, m_2) which occur in any relation of $\mathfrak{M}$, occur also in $R_j^{\mathfrak{M}}$ or in $\left(R_j^{\mathfrak{M}}\right)^{-1}$. Additional unary relation symbols $U_{B,\mathrm{self}}$ encode self loops (m, m) in $B^{\mathfrak{M}}$.

Let $\mathbf{s}_{\mathcal{C}} = \langle \varphi, \psi_C : C \in \mathcal{C} \rangle$ is the translation scheme given as follows.

- $\varphi(x) = (x \approx x)$.
- For every unary relation symbol $U \in \mathcal{C}$, $\psi_U(x) = U(x)$.
- For every binary relation symbol $B \in \mathcal{C}$, $B^{\mathbf{s}_{\mathcal{C}}^{\star}(\mathfrak{T})}$ consists of all pairs (v, u) such that (a) $(v, u) \in R_j^{\mathfrak{T}}$ and $v \in U_{B,j}^{\mathfrak{T}}$ or (b) $(u, v) \in R_j^{\mathfrak{T}}$ and $u \in U_{B,\mathrm{inv},j}^{\mathfrak{T}}$. This is given by

$$\begin{aligned}\psi_B(x,y) \quad &= \quad \neg(x \approx y) \wedge \bigvee_{j \in [k]} \left((R_j(x,y) \wedge U_{B,j}(x)) \vee (R_j(y,x) \wedge U_{B,\mathrm{inv},j}(y)) \right) \\ & \qquad \vee (x \approx y) \wedge U_{B,\mathrm{self}}(x)\end{aligned}$$

Let $\mathfrak{M}$ be a $\mathcal{C}$-structure with universe M. Let $\mathfrak{A}_0$ be an aligned k-tree such that $\mathrm{Gaif}(\mathfrak{M}) = \mathrm{Gaif}(\mathfrak{A}_0)$ guaranteed by Lemma 17. Let $\mathfrak{A}_1$ be the expansion of $\mathfrak{A}_0$ such that, for every $B \in \mathcal{C}$ and $j \in [k]$,

$$\begin{aligned} U_{B,j}^{\mathfrak{A}_1} &= \left\{ m_1 \mid (m_1, m_2) \in R_j^{\mathfrak{A}_1} \cap B^{\mathfrak{M}} \right\} \\ U_{B,\mathrm{inv},j}^{\mathfrak{A}_1} &= \left\{ m_1 \mid (m_2, m_1) \in \left(R_j^{A_1}\right)^{-1} \cap B^{\mathfrak{M}} \right\} \\ U_{B,\mathrm{self}}^{\mathfrak{A}_1} &= \{ m \mid (m, m) \in B^{\mathfrak{M}} \} \end{aligned}$$

Self loops are encoded by the relations of the form $U_{B,\mathrm{self}}^{\mathfrak{A}_1}$. Consider $a, b \in M$ which are distinct. We divide into cases.

- Assume (a, b) is not an edge of $\mathrm{Gaif}(\mathfrak{M})$. By $\mathrm{Gaif}(\mathfrak{M}) = \mathrm{Gaif}(\mathfrak{A}_1)$, neither (a, b) nor (b, a) belong to any $R_j^{\mathfrak{T}_0} = R_j^{\mathfrak{T}_1}$. Hence, $\mathfrak{A}_1 \not\models \psi_B(a, b)$, i.e. $(a, b) \notin B^{\mathbf{s}_{\mathcal{C}}^{\star}(\mathfrak{A}_1)}$, for any binary relation symbol $B \in \mathcal{C}$.
- Assume (a, b) is an edge of $\mathrm{Gaif}(\mathfrak{M})$. By $\mathrm{Gaif}(\mathfrak{M}) = \mathrm{Gaif}(\mathfrak{A}_0)$, either (a, b) or (b, a) belong to some $R_j^{\mathfrak{A}_0} = R_j^{\mathfrak{A}_1}$. We have: $(a, b) \in B^{\mathfrak{M}}$ iff exists $j \in [k]$ such that $[(a, b) \in R_j^{\mathfrak{A}_1}$ and $a \in U_{B,j}^{\mathfrak{A}_1}]$ or $[(b, a) \in R_j^{\mathfrak{A}_1}$ and $b \in U_{B,\mathrm{inv},j}^{\mathfrak{A}_1}]$ iff $\mathfrak{A}_1 \models \psi_B(a, b)$ iff $(a, b) \in B^{\mathbf{s}_{\mathcal{C}}^{\star}(\mathfrak{A}_1)}$.

In both cases $(a, b) \in B^{\mathfrak{M}}$ iff $(a, b) \in B^{\mathbf{s}_{\mathcal{C}}^{\star}(\mathfrak{A}_1)}$. We get that $\mathfrak{M} = \mathbf{s}_{\mathcal{C}}^{\star}(\mathfrak{A}_1)$ and (1) follows.

Now we turn to (2). By Lemma 17(1), the Gaifman graph of $\mathfrak{A}$ is a partial $(k-1)$-tree. By definition of ψ_B, for every binary $B \in \mathcal{C}$, $B^{\mathbf{s}_{\mathcal{C}}^{\star}(\mathfrak{A})} \subseteq \bigcup_{j \in [k]} R_j^{\mathfrak{A}} \cup \left(R_j^{\mathfrak{A}}\right)^{-1}$, implying that $\mathrm{Gaif}(\mathbf{s}_{\mathcal{C}}^{\star}(\mathfrak{A}))$ is a partial $(k-1)$-tree.

The translation scheme $\mathbf{tr}_{\mathcal{C}}$: from structures of bounded tree-width to labeled trees. Let $\mathbf{i}_{\mathcal{C}}$ and $\mathbf{u}_{\mathcal{C}}$ be the translation schemes which extend $\mathbf{i}$ and $\mathbf{u}$ respectively as follows. The unary relations C of $\mathcal{N}_{\mathcal{C}}$ are additionally defined under $\mathbf{i}_{\mathcal{C}}$ and $\mathbf{u}_{\mathcal{C}}$ to be $\psi_C(x) = C(x)$. Hence, $\mathbf{i}_{\mathcal{C}}$ is a translation scheme for $\mathcal{R} \cup \mathcal{V} \cup \mathcal{N}_{\mathcal{C}}$ over $\mathcal{V} \cup \mathcal{N}_{\mathcal{C}}$, $\mathbf{u}_{\mathcal{C}}$ is a translation scheme for $\mathcal{R} \cup \mathcal{N}_{\mathcal{C}}$ over $\mathcal{R} \cup \mathcal{V} \cup \mathcal{N}_{\mathcal{C}}$, and $\mathbf{s}_{\mathcal{C}}$ is a translation scheme for $\mathcal{C}$ over $\mathcal{R} \cup \mathcal{N}_{\mathcal{C}}$.

Before we define $\mathbf{tr}_{\mathcal{C}}$, we need to introduce the notion of composition of translation schemes. Let $\mathcal{C}_0$, $\mathcal{C}_1$ and $\mathcal{C}_2$ be vocabularies. Let $\mathbf{t}_i = \left\langle \phi^i, \psi^i_C : C \in \mathcal{C}_i \right\rangle$ be a translation scheme for $\mathcal{C}_i$ over $\mathcal{C}_{i-1}$. The **composition** of $\mathbf{t}_1$ and $\mathbf{t}_2$, denoted $\mathbf{t}_1 \circ \mathbf{t}_2$, is the translation scheme given by $\mathbf{t} = \langle \phi, \psi_C : C \in \mathcal{C}_i \rangle$ such that

1. $\phi(x) = (\mathbf{t}^1)^\sharp((\mathbf{t}^2)^\sharp(x \approx x))$
2. For every $C \in \mathrm{un}(\mathcal{C})$, $\psi_C(x) = (\mathbf{t}^1)^\sharp((\mathbf{t}^2)^\sharp(C(x)))$
3. For every $C \in \mathrm{bin}(\mathcal{C})$, $\psi_C = (\mathbf{t}^1)^\sharp((\mathbf{t}^2)^\sharp(C(x,y)))$

The translation scheme $\mathbf{t}_{\mathcal{C}}$ is the composition of $\mathbf{i}_{\mathcal{C}}$, $\mathbf{u}_{\mathcal{C}}$ and $\mathbf{s}_{\mathcal{C}}$, i.e. $\mathbf{t}_{\mathcal{C}} = \mathbf{i}_{\mathcal{C}} \circ (\mathbf{u}_{\mathcal{C}} \circ \mathbf{s}_{\mathcal{C}})$. $\mathbf{t}_{\mathcal{C}}$ is a translation scheme for $\mathcal{C}$ over $\mathcal{V} \cup \mathcal{N}_{\mathcal{C}}$. As a consequence of Lemma 18, we have:

> **Lemma 20.** *There is a vocabulary $\mathcal{N}_{\mathcal{C}}$ consisting of unary relation symbols only, such that for every $\mathcal{C}$-structure $\mathfrak{M}$, $\mathfrak{M}$ has tree-width at most k iff there is an $\mathcal{V} \cup \mathcal{N}_{\mathcal{C}}$-structure $\mathfrak{A}$ such that $\mathbf{t}^\star_{\mathcal{C}}(\mathfrak{A}) = \mathfrak{M}$ and $\mathfrak{A}$ is an aligned k-tree encoding.*

Proof of Lemma 14. Let $\mathcal{C}_{bnd}$ and $\mathcal{C}_{unb}$ be vocabularies such that $\mathcal{C}_{bnd} \cap \mathcal{C}_{unb}$ only contains unary relation symbols. Let $\mathcal{D}_{bnd} = \mathcal{V} \cup \mathcal{N}_{\mathcal{C}_{bnd}}$. We define $\mathbf{tr}$ which extends $\mathbf{t}_{\mathcal{C}_{bnd}}$ as follows: For every $C \in \mathcal{C}_{unb} \backslash \mathcal{C}_{bnd}$, let $\psi_C(x) = C(x)$ or $\psi_C(x,y) = C(x,y)$. I.e. $\mathbf{tr}$ copies the relations of $\mathcal{C}_{unb} \backslash \mathcal{C}_{bnd}$ from the source structure to the target structure without change (except taking the projection to the universe of the target structure imposed by the formula $\phi(x)$). Let *dom* = enc from Lemma 16.

(a) ϕ is quantifier-free since $\mathbf{t}_{\mathcal{C}}$ is obtained from the composition of translation schemes in which ϕ is quantifier-free.
(b) ψ_C is quantifier-free for every $C \in \mathcal{C}_{unb} \backslash \mathcal{C}_{bnd}$ by the definition in this subsection. ψ_C is quantifier-free for every $C \in \mathrm{un}(\mathcal{C}_{bnd})$ since we have in fact $\psi_C(x) = C(x)$; this is true because it is true for each of $\mathbf{i}_{\mathcal{C}}$, $\mathbf{u}_{\mathcal{C}}$ and $\mathbf{s}_{\mathcal{C}}$.
(c) For every $C \in \mathcal{C}_{bnd}$, ψ_C is defined in the translation scheme $\mathbf{tr}_{\mathcal{C}_{bnd}}$. Since $\mathbf{tr}_{\mathcal{C}_{bnd}}$ is a translation scheme for $\mathcal{C}_{bnd}$ over $\mathcal{V} \cup \mathcal{N}_{\mathcal{C}_{bnd}} = \mathcal{D}_{bnd}$, $\psi_C \in \mathrm{MSO}(\mathcal{D}_{bnd})$.
(d) This item follows from Lemma 20 and using that *dom* = enc defines the class of aligned k-tree encodings on structures $\mathfrak{A}$ in which $s^{\mathfrak{A}}$ is a binary tree.

8.3. **Proof of Lemma 15.** For a leaf b, $\mathrm{hin}_q(\mathfrak{T}_b)$ depends only on the unary relations which b satisfies. By Theorem 5, for a vertex b with one child b_0 (two children b_0, b_1), $\mathrm{hin}_q(\mathfrak{T}_b)$ depends only on the unary relations which b satisfies and on $\mathrm{hin}_q(\mathfrak{T}_{b_0})$ (on $\mathrm{hin}_q(\mathfrak{T}_{b_0})$ and $\mathrm{hin}_q(\mathfrak{T}_{b_1})$).

Let

$$\Theta^{\mathrm{hin}}_{\mathcal{C},q} = \mathrm{part} \wedge \mathrm{leaves} \wedge \mathrm{ints}_1 \wedge \mathrm{ints}_2$$

where part says that $\{C_\epsilon : \epsilon \in \mathscr{HIN}_{\mathcal{C},q}\}$ partition the universe, and leaves, ints_1, and ints_2 define the C_ϵ for the leaves respectively the internal vertices of $\mathfrak{T}_0$ with one or two children.

We give part, leaves, ints_1 and ints_2 below. There are C^2 formulas $\mathrm{leaf}(x)$, $\mathrm{int}_1(x)$, and $\mathrm{int}_2(x)$, which express that x is a leaf, has one child, or has two children, respectively. Let

$$\mathrm{part} = \forall x \left(\left(\bigvee_{\epsilon \in \mathscr{HIN}_{\mathcal{C},q}} C_\epsilon(x) \right) \wedge \left(\bigwedge_{\epsilon_1 \neq \epsilon_2 \in \mathscr{HIN}_{\mathcal{C},q}} (\neg C_{\epsilon_1}(x) \vee \neg C_{\epsilon_2}(x)) \right) \right).$$

For $\mathcal{U} \subseteq \mathrm{un}(\mathcal{C})$, let $\mathfrak{O}_\mathcal{U}$ be a $\mathcal{C}$-structure with universe O of size 1 satisfying that $U^{\mathfrak{O}_\mathcal{U}} = \emptyset$ iff $U \in \mathcal{U}$, and $\mathrm{root}^{\mathfrak{O}_\mathcal{U}} = O$. Let

$$\begin{aligned}
\mathrm{ints}_1 &= \forall x \left(\mathrm{int}_1(x) \to \bigwedge_{\mathcal{U},\epsilon_1,\epsilon_2} ((\mathrm{this}_\mathcal{U}(x) \wedge \mathrm{child}_{\epsilon_2}(x)) \to C_{\epsilon_1}(x)) \right) \\
\mathrm{this}_\mathcal{U}(x) &= \bigwedge_{U \in \mathrm{un}(\mathcal{C})} \neg^{(U \in \mathcal{U})} U(x) \\
\mathrm{child}_{\epsilon_2}(y) &= \exists y \, s(x,y) \wedge C_{\epsilon_2}(y)
\end{aligned}$$

$\bigwedge_{\mathcal{U},\epsilon_1,\epsilon_2}$ ranges over $\epsilon_1, \epsilon_2 \in \mathscr{HIN}_{\mathcal{C},q}$ and $\mathcal{U} \subseteq \mathrm{un}(\mathcal{C})$ such that for every structure $\mathfrak{A} \in \mathscr{K}_{\mathrm{root}}$ with $\mathrm{hin}(\mathfrak{A}) = \epsilon_2$, $\mathrm{hin}(\mathfrak{O}_\mathcal{U} \overset{\rightarrow}{\circ} \mathfrak{A}) = \epsilon_1$, The notation $\neg^X Y$ stands for Y if X is false, and for $\neg Y$ otherwise.

We define

$$\mathrm{leaves} = \forall x \left(\mathrm{leaf}(x) \to \bigwedge_{\mathcal{U},\epsilon_1} \mathrm{this}_\mathcal{U}(x) \to C_{\epsilon_1}(x) \right)$$

$\bigwedge_{\mathcal{U},\epsilon_1}$ ranges over $\epsilon_1 \in \mathscr{HIN}_{\mathcal{C},q}$ and $\mathcal{U} \subseteq \mathrm{un}(\mathcal{C})$, where $\mathrm{hin}(\mathfrak{O}_\mathcal{U}) = \epsilon_1$.

Finally, ints_2 is defined as follows:

$$\begin{aligned}
\mathrm{ints}_2 &= \forall x (\mathrm{int}_2(x) \to (\mathrm{intdist}(x) \wedge \mathrm{intsame})) \\
\mathrm{intdist}(x) &= \bigwedge_{\mathcal{U},\epsilon_1,\epsilon_2,\epsilon_3} ((\mathrm{this}_\mathcal{U}(x) \wedge \mathrm{child}_{\epsilon_2}(x) \wedge \mathrm{child}_{\epsilon_3}(x)) \to C_{\epsilon_1}(x)) \\
\mathrm{intsame}(x) &= \bigwedge_{\mathcal{U},\epsilon_1,\epsilon_2} ((\mathrm{this}_\mathcal{U}(x) \wedge \mathrm{children}_{\epsilon_2}(x)) \to C_{\epsilon_1}(x)) \\
\mathrm{children}_{\epsilon_2}(x) &= \exists^{=2} y \, (s(x,y) \wedge C_{\epsilon_2}(y))
\end{aligned}$$

$\bigwedge_{\mathcal{U},\epsilon_1,\epsilon_2,\epsilon_3}$ in $\mathrm{intdist}(x)$ ranges over $\epsilon_1, \epsilon_2, \epsilon_3 \in \mathscr{HIN}_{\mathcal{C},q}$ and $\mathcal{U} \subseteq \mathrm{un}(\mathcal{C})$ such that $\epsilon_2 \neq \epsilon_3$, and for every two structure $\mathfrak{A}, \mathfrak{B} \in \mathscr{K}_{\mathrm{root}}$ with $\mathrm{hin}(\mathfrak{A}) = \epsilon_2$ and $\mathrm{hin}(\mathfrak{B}) = \epsilon_3$, $\mathrm{hin}((\mathfrak{O}_\mathcal{U} \overset{\rightarrow}{\circ} \mathfrak{A}) \overset{\rightarrow}{\circ} \mathfrak{B}) = \epsilon_1$. $\bigwedge_{\mathcal{U},\epsilon_1,\epsilon_2}$ in $\mathrm{intsame}(x)$ ranges over $\epsilon_1, \epsilon_2 \in \mathscr{HIN}_{\mathcal{C},q}$ and $\mathcal{U} \subseteq \mathrm{un}(\mathcal{C})$ such that for every structure $\mathfrak{A} \in \mathscr{K}_{\mathrm{root}}$ with $\mathrm{hin}(\mathfrak{A}) = \epsilon_2$, $\mathrm{hin}(\mathfrak{O}_\mathcal{U} \overset{\rightarrow}{\circ} \mathfrak{A}) = \epsilon_1$.

By definition, $\Theta^{\mathrm{hin}}_{\mathcal{C},q} \in C^2$. $\mathfrak{T}_0$ is a $\mathcal{C}$-structure in which s is interpreted as a binary tree. Let $\mathfrak{T}_1$ be the expansion of $\mathfrak{T}_0$ such that, for every element b of the universe $T_1 = T_0$ of $\mathfrak{T}_1$, $b \in C^{T_1}_{\mathrm{hin}(\mathfrak{T}_b)}$, where $\mathfrak{T}_b$ is the subtree of $\mathfrak{T}_0$ rooted at b. In particular, for the root r of $\mathfrak{T}_0$, $r \in C^{T_1}_{\mathrm{hin}(\mathfrak{T}_r)} = C^{T_1}_{\mathrm{hin}(\mathfrak{T}_0)}$. We have $\mathfrak{T}_1 \models \Theta^{\mathrm{hin}}_{\mathcal{C},q}$, and hence (i) holds. Since $\mathfrak{T}_1$ is the only expansion of $\mathfrak{T}_0$ which satisfies $\Theta^{\mathrm{hin}}_{\mathcal{C},q}$, (ii) holds.

8.4. **Proof of Lemma 4.** We assume $2\text{-}\mathsf{tp}(\mathfrak{M}_1) = 2\text{-}\mathsf{tp}(\mathfrak{M}_2)$. Let $u_1, v_1 \in M$ be two elements and let $\lambda = 2\text{-}\mathsf{tp}^{\mathfrak{M}_1}(u_1, v_1)$. Then there are two elements $u_2, v_2 \in M$ such that $\lambda = 2\text{-}\mathsf{tp}^{\mathfrak{M}_2}(u_2, v_2)$ because of $2\text{-}\mathsf{tp}(\mathfrak{M}_1) = 2\text{-}\mathsf{tp}(\mathfrak{M}_2)$. Thus, $\mathfrak{M}_1 \models \chi(u_1, v_1)$ implies $\mathfrak{M}_2 \models \chi(u_2, v_2)$. Because u_1, v_1 have been chosen arbitrarily, we get that $\mathfrak{M}_1 \models \phi$ implies $\mathfrak{M}_2 \models \phi$. The other direction is symmetric.

8.5. **Proof of Lemma 6.**

Proof. Let $z = 2|\mathcal{T}|^2 + 1$. We set $\mathrm{colors}(\mathcal{T}) = \{A_i \mid 1 \leq i \leq z\}$, where the A_i are fresh unary relation symbols.

Let $\mathfrak{M}$ be a $\mathcal{C}_{unb}$-structure with universe M. By Lemma 7, there is a proper z-coloring of $G_{\mathcal{T}}^{\mathfrak{M}}$. We expand $\mathfrak{M}$ to a $(\mathcal{C}_{unb} \cup \text{colors}(\mathcal{T}))$-structure $\mathfrak{L}$ as follows: for all elements $u \in M$ we have $A_i^{\mathfrak{L}}(u)$ iff u has color i in the proper z-coloring of $G_{\mathcal{T}}^{\mathfrak{M}}$. Clearly, $\mathfrak{L}$ is a chromatic $(\mathcal{C}_{unb} \cup \text{colors}(\mathcal{T}))$-structure. $\square$

8.6. **Proof of Lemma 7.** We proceed by induction on the number of vertices $n = |V|$. The case $n = 1$ is trivial. Assume that $n > 1$ and that the lemma holds for $n-1$. We have $\sum_{v\in V} deg(v) = 2 \cdot \sum_{v \in V} deg^+(v) \leq 2nk$. Thus there is a vertex $v \in V$ with $deg(v) \leq 2k$. By the induction hypothesis, the graph $G{-}v$ has a proper $2k+1$-coloring. Because of $deg(v) \leq 2k$ the proper $2k+1$-coloring of $G{-}v$ can be extended to a proper $2k+1$-coloring of G.

8.7. **Proof of Lemma 10.** Let $v \in M$ be an element with $2\text{-}\mathfrak{tp}^{\mathfrak{L}|\varepsilon}(u,v) = \lambda$. We get $\mathfrak{L} \models P_\lambda(u)$ because of $\mathfrak{L} \models \zeta_{unb}$. Because of $1\text{-}\mathfrak{tp}^{\mathfrak{L}}(u) = 1\text{-}\mathfrak{tp}^{\mathfrak{L}'}(u)$ we have $\mathfrak{L}' \models P_\lambda(u)$. Because of $\mathfrak{L}' \models \zeta_{bnd}$ we have $\lambda = 2\text{-}\mathfrak{tp}^{\mathfrak{L}'|\varepsilon}(u,w)$ for some $w \in M$. $rank_u^i(\mathfrak{L},\mathfrak{L}') = 0$ implies that $\mathfrak{L}' \models R_i(u,v)$. Because of $\mathfrak{L}' \models \delta$ we get $v = w$. In the same way one can show that $2\text{-}\mathfrak{tp}^{\mathfrak{L}'|\varepsilon}(u,v) = \lambda$ implies $2\text{-}\mathfrak{tp}^{\mathfrak{L}|\varepsilon}(u,v) = \lambda$.

8.8. **Proof of Lemma 9.** We show $\mathfrak{L}|_{\mathcal{C}_{bnd}} = \mathfrak{L}'|_{\mathcal{C}_{bnd}}$. Let $u, v \in M$ with $\mathfrak{L} \models B(u,v)$ for some $B \in \text{bin}(\mathcal{C}_{bnd})$. Because of $\mathfrak{L} \models \theta$ we have $\mathfrak{L} \models R_i(u,v)$ or $\mathfrak{L} \models R_i(v,u)$ for some $i \in [k]$. Because of $rank(\mathfrak{L},\mathfrak{L}') = 0$ we have $rank_u^i(\mathfrak{L},\mathfrak{L}') = 0$. By Lemma 10 we have $2\text{-}\mathfrak{tp}^{\mathfrak{L}|\varepsilon}(u,v) = 2\text{-}\mathfrak{tp}^{\mathfrak{L}'|\varepsilon}(u,v)$. Thus, $\mathfrak{L}' \models B(u,v)$. In the same way one can show that $\mathfrak{L}' \models B(u,v)$ implies $\mathfrak{L} \models B(u,v)$. Thus, $\mathfrak{L}|_{\mathcal{C}_{bnd}} = \mathfrak{L}'|_{\mathcal{C}_{bnd}}$.

We show $\text{Gaif}(\mathfrak{L}|_{\mathcal{C}_{bnd}}) = \text{Gaif}(\mathfrak{L}')$: We have that $\text{Gaif}(\mathfrak{L}') = \text{Gaif}(\mathfrak{L}'|_{\mathcal{R}})$ because of $\mathfrak{L}' \models \theta$ and $\mathfrak{L}' \models \epsilon$. We have $\text{Gaif}(\mathfrak{L}|_{\mathcal{C}_{bnd}}) = \text{Gaif}(\mathfrak{L}|_{\mathcal{R}})$ because of $\mathfrak{L} \models \theta$. We have $\text{Gaif}(\mathfrak{L}'|_{\mathcal{R}}) = \text{Gaif}(\mathfrak{L}|_{\mathcal{R}})$ because of $rank(\mathfrak{L},\mathfrak{L}') = 0$. Thus, the claim follows.

8.9. **The induced translation $\mathbf{t}^\sharp$.** We spell out the inductive definition of the induced translation.

Let $\mathcal{C}_0$ and $\mathcal{C}_1$ be vocabularies. Given a translation scheme $\mathbf{t} =< \phi, \psi_C : C \in \mathcal{C}_0 >$ for $\mathcal{C}_0$ over $\mathcal{C}_1$ we define the induced translation $\mathbf{t}^\sharp$ to be a function from MSO($\mathcal{C}_0$)-formulas to $\mathcal{C}_1$-formulas inductively as follows:

1. For $C \in \text{un}(\mathcal{C}_0)$ or for monadic second order variables C, and for $\theta = C(x)$, we put
$$\mathbf{t}^\sharp(\theta) = \psi_C(x) \wedge \phi(x)$$
2. For $C \in \text{bin}(\mathcal{C}_0)$ and $\theta = C(x,y)$, we put
$$\mathbf{t}^\sharp(\theta) = \psi_C(x,y) \wedge \phi(x) \wedge \phi(y)$$
3. For $x \approx y$, we put
$$\mathbf{t}^\sharp(\theta) = x \approx \wedge \phi(x) \wedge \phi(y)$$
4. For the Boolean connectives the translation distributes, i.e.
    - if $\theta = \theta_1 \vee \theta_2$ then
    $$t^\sharp(\theta) = (\mathbf{t}^\sharp(\theta_1) \vee \mathbf{t}^\sharp(\theta_2))$$
    - if $\theta = \neg\theta_1$ then
    $$t^\sharp(\theta) = \neg \mathbf{t}^\sharp(\theta_1)$$

(5) For the existential quantifiers, we relativize to ϕ:
If $\theta = \exists y \, \theta_1$, we put

$$\mathbf{t}^\sharp(\theta) = \exists y \, (\phi(y) \wedge \mathbf{t}^\sharp(\theta_1))$$

If $\theta = \exists U \, \theta_1$, we put

$$\mathbf{t}^\sharp(\theta) = \exists U \, (\mathbf{t}^\sharp(\theta_1) \wedge \forall y \, U(y) \to \phi(y))$$

We have somewhat simplified the presentation in [19, Definition 2.3] to fit our setting.

TU Vienna
E-mail address: {`kotek,zuleger@forsyte.at`}